

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 February 2003 (27.02.2003)

PCT

(10) International Publication Number
WO 03/015776 A1

(51) International Patent Classification: **A61K 31/426**,
31/427, 31/506, 31/497, C07D 417/04, 417/14, A61P
31/12, 37/02, 37/08, 29/00, C07D 277/24

(74) Common Representative: JANSSEN PHARMACEU-
TICA N.V.; Lies Vervoort, Johnson & Johnson Patent Law
Department, Turnhoutseweg 30, B-2340 Beerse (BE).

(21) International Application Number: PCT/EP02/08956

(22) International Filing Date: 9 August 2002 (09.08.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
01203088.8 13 August 2001 (13.08.2001) EP

(71) Applicant (for all designated States except US):
JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turn-
houtseweg 30, B-2340 Beerse (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **LOVE, Christo-
pher, John** [GB/BE]; Janssen Pharmaceutica N.V.,
Turnhoutseweg 30, B-2340 Beerse (BE). **VAN WAUWE,
Jean, Pierre, Frans** [BE/BE]; Janssen Pharmaceutica
N.V., Turnhoutseweg 30, B-2340 Beerse (BE). **DE BRA-
BANDER, Marc, J.** [BE/BE]; Valerijalaan 2, B-2980
Zoersel (BE). **MOSES, Roger, Clive** [GB/GB]; 18
Femleigh Road, Walebridge, Cornwall PL27 7AX (GB).
GONCHARENKO, Mykhal'yo [UA/CA]; 875 Wonder-
land Road South, Apt. #1202, London, Ontario N6K 3N1
(CA). **COOYMANS, Ludwig, Paul** [BE/BE]; Janssen
Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse
(BE). **VANDERMAESEN, Nele** [BE/BE]; Janssen Phar-
maceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE).
DIELS, Gaston, Stanislas, Marcella [BE/BE]; Janssen
Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse
(BE). **SIBLEY, Anthony, William** [GB/GB]; 1 Camelot
View, Camelford, Cornwall PL32 9TU (GB). **NOULA,
Caterina** [GR/GB]; 3 Beech Drive, St. Columb Major,
Cornwall TR9 6SZ (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii)) for the following designations AE,
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES,
FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG,
MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU,
SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG,
UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS,
MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB,
GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF,
BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,
TD, TG)
- as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for all designations
- of inventorship (Rule 4.17(iv)) for US only

[Continued on next page]

WO 03/015776 A1

(54) Title: 2,4,5-TRISUBSTITUTED THIAZOLYL DERIVATIVES AND THEIR ANTIINFLAMMATORY ACTIVITY



(57) Abstract: This invention concerns the use of a compound of formula (I), a N-oxide, a phar-
maceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form
thereof, wherein Z is halo; C₁₋₆alkyl; C₁₋₆alkyl substituted with hydroxy, carboxyl, cyano, amino,
mono -or di(C₁₋₆alkyl)amino, aminocarbonyl, mono -or di(C₁₋₆alkyl)aminocarbonyl, C₁₋₆alkyloxy-
carbonyl or C₁₋₆alkyloxy; polyhaloC₁₋₄alkyl; C₁₋₄alkyloxy; cyano; amino; aminocarbonyl; mono
-or di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; H₂N-S(=O)₂-; mono -or di(C₁₋₆alkyl)amino-S(=O)₂;
-C(=N-R¹)NR²R²; Q is an optionally substituted carbocyclic or an optionally substituted heterocycle; L is substituted phenyl or an
optionally substituted monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a bicyclic partially saturated or
aromatic heterocycle; aryl is optionally substituted phenyl; for the manufacture of a medicament for the prevention or the treatment
of diseases mediated through TNF-α and/or IL-12.



Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 03/015776 A1

2,4,5-TRISUBSTITUED THIAZOLYL DERIVATIVES AND THEIR ANTIINFLAMMATORY ACTIVITY

The present invention is concerned with 2,4,5-trisubstituted thiazolyl derivatives having proinflammatory cytokine production inhibiting properties. The invention further relates to methods for their preparation and pharmaceutical compositions comprising them. The invention also relates to the use of 2,4,5-trisubstituted thiazolyl derivatives for the manufacture of a medicament for the prevention or the treatment of diseases mediated through TNF- α and/or IL-12.

10

WO 00/35911 describes acetal derivatives as TNF- α inhibitors.

WO 96/03392 describes sulphonyl derivatives for treating inflammation.

WO 98/01449 describes pyrimidine fused compounds as antiallergic and antiinflammatory agents.

15 US 5,240,929 describes 2-heterocyclic-5-hydroxy-1,3-pyrimidines useful as antiinflammatory agents.

Z. Chem., 1969, 9(5), 186-187 describes thiazolyl quinolines as fluorescence indicators.

20 EP 117,082 describes thiazole derivatives as cardiogenic, blood pressure regulating and anti-ulcer agents.

Chem.Pharm. Bull., 1982, 30(6), 1974-1979 discloses studies on tertiary amine oxides.

WO 97/05131 describes heteroarylcarboxamides as agrochemical and medical fungicides.

25 JP 91-144612 relates to isoxazoles as disinfectants, antiseptics, anti-inflammatory agents, bactericides, viricides.

Arch.Pharm. 1981, 314(9), 744-750 describes the synthesis and antibacterial effect of 2-aryl-4-R-5 glyoxyl-thiazoles.

Z.Naturforsch. B Chem. Sci. 1990, 45(12), 1695-1708 concerns cycloaddition reactions with 5-azido-4-(trifluoromethyl)-1,3-azoles.

30 Synthesis 1988, 3, 194-198 describes methods for regioselective introduction of trifluoromethyl groups into heteroarenes.

J.Med.Chem. 1988, 31 (6), 1197-1204 describes the synthesis of oxazole derivatives as hypolipidemic, anticholesteremic and blood platelet aggregation inhibiting agents.

Chem.Ber. 1982, 115(7), 2494-2507 relates to the synthesis of 1,3-azoles.

35 DD 258165 describes quinoxaline derivatives as herbicides and fungicides.

Z.Chem. 1979, 19(1), 21-22 relates to heterocyclic substituted thiazoles as pesticides.

-2-

Indian J.Chem., Sect. B, 1976, 14B(7), 552-555 concerns the synthesis and anti-inflammatory activity of bithiazolyl derivatives.

Justus Liebigs Ann. Chem. 1974, 8, 1195-1205 describes the synthesis of thiazole compounds.

5 DE 1959307 concerns benzoxazole derivatives as fluorescent whiteners.

GB 1189008 relates to benzoxazole derivatives as fluorescent whiteners.

JP 41012946 describes benzoxazole and benzimidazole derivatives as optical brightening organic fibers.

Chem.Ber. 1967, 100(7), 2184-2187 describes the synthesis of thiazole derivatives.

10 US 6,231,786 describes fluorinated azoles and their use in liquid crystalline mixtures.


WO 01/30778 discloses thiazoles and imidazopyridines for treating TNF and IL-1 mediated diseases.

WO 98/08830 and WO 98/08841 disclose thiazole derivatives having PDE IV inhibiting properties.

15 WO 01/64674 describes 2,4-disubstituted thiazolyl derivatives as TNF- α and/or IL-12 inhibitors.

20 The compounds of the present invention are distinguishable from the prior art because of their structure, pharmacological activity, potency or physicochemical characteristics (improved chemical stability, improved solubility).

25 The present invention relates to the use of a compound for the manufacture of a medicament for the prevention or the treatment of diseases mediated through TNF- α (Tumor Necrosis Factor-alpha) and/or IL-12 (Interleukin 12) wherein the compound is a compound of formula



a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof,
wherein

30 Z is halo; C₁₋₆alkyl; C₁₋₆alkyl substituted with hydroxy, carboxyl, cyano, amino, mono -or di(C₁₋₆alkyl)amino, aminocarbonyl, mono -or di(C₁₋₆alkyl)aminocarbonyl, C₁₋₆alkyloxycarbonyl or C₁₋₆alkyloxy; polyhaloC₁₋₄alkyl; C₁₋₄alkyloxy; cyano; amino; aminocarbonyl; mono -or di(C₁₋₆alkyl)aminocarbonyl;

-3-


C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; H₂N-S(=O)₂-; mono -or
 di(C₁₋₆alkyl)amino-S(=O)₂; -C(=N-R^x)NR^yR^z;

R^x is hydrogen, C₁₋₆alkyl, cyano, nitro or -S(=O)₂-NH₂;

R^y is hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl or C₂₋₆alkynyl;

5 R^z is hydrogen or C₁₋₆alkyl;

Q is C₃₋₆cycloalkyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl,
 pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, phenyl,
 pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl,
 benzimidazolyl, indazolyl, or imidazopyridyl, wherein each of said ring systems
 10 may optionally be substituted with up to three substituents each of said
 substituents independently being selected from halo; hydroxy; cyano; carboxyl;
 azido; amino; mono- or di(C₁₋₆alkyl)amino; C₁₋₆alkylcarbonylamino; C₁₋₆alkyl;
 C₂₋₆alkenyl; C₂₋₆alkynyl; C₃₋₆cycloalkyl; C₁₋₆alkyl substituted with hydroxy,
 C₁₋₆alkyloxy, amino, mono- or di(C₁₋₄alkyl)amino; C₁₋₆alkyloxy; C₁₋₆alkylthio;
 15 C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; arylC₁₋₆alkyloxy; aryloxy;
 polyhaloC₁₋₆alkyl; polyhaloC₁₋₆alkyloxy; polyhaloC₁₋₆alkylcarbonyl; C₁₋₄alkyl-
 S(=O)_n- or R¹HN-S(=O)_n-, with R¹ representing hydrogen, or a radical of formula




with A being O, S or a bivalent radical of formula -CR²=N- with CR²
 20 attached to N of formula (a-1); and

R² being hydrogen, C₁₋₆alkyl or C₁₋₆alkyloxy;

or

Q is a radical of formula



25 wherein B₁ and B₂ each independently are O, NR³, CH₂ or S, with R³ being
 hydrogen or C₁₋₄alkyl;

B₃ is O or NR⁴ with R⁴ being hydrogen or C₁₋₄alkyl;

q is an integer with value 1 to 4;

r is an integer with value 1 to 3;

30 n is an integer with value 1 or 2;

L is phenyl substituted with up to 4 substituents each substituent independently being
 selected from C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; aminocarbonyl; mono-
 or di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyl-C(=O)-NH-; C₁₋₆alkyloxy-C(=O)-NH-;

H₂N-C(=O)-NH-; mono- or di(C₁₋₄alkyl)amino-C(=O)-NH-; Het-NH-; Het¹-NH-;
 -NH-C(=N-R^x)NR^yR^z; -C(=N-R^x)NR^yR^z; Het¹; or a radical of formula
 -X-C₁-Y₁-C₂-Y₂-C₃-Y₃-C₄-Z (c-1) wherein

X represents NR⁵, O, S or a direct bond;

5 C₁, C₂, C₃ and C₄ each independently represent C₁₋₆alkanediyl,
 C₂₋₆alkenediyl, C₂₋₆alkynediyl or a direct bond;

Y₁, Y₂ and Y₃ each independently represent NR⁵, O, S or a direct bond;

Z is hydrogen, halo, cyano, hydroxy, carboxyl, -P(=O)(OH)H,

10 -P(=O)(OH)₂, -P(=O)(OH)CH₃, -P(=O)(OH)(OCH₃),
 -P(=O)(OH)(OCH₂CH₃), -P(=O)(OH)NH₂, -S(=O)₂H, -S(=O)₂(OH),
 -S(=O)₂NH, -C(=O)-NH-S(=O)₂-H, tetrazolyl, 3-hydroxy-isothiazolyl,
 3-hydroxy-isoxazolyl, 3-hydroxy-thiadiazolyl, mercaptotetrazolyl,
 3-mercapto-triazolyl, 3-sulfinyl-triazolyl, 3-sulfonyl-triazolyl;

R⁵ is hydrogen, C₁₋₆alkyl or -C(=NH)-N(R^z)₂; and wherein

15 from 1 to 3 hydrogen atoms of the C₁₋₆alkyl, C₁₋₆alkanediyl,
 C₂₋₆alkenediyl or C₂₋₆alkynediyl groups in the definitions of R⁵ and the
 radical of formula (c-1) may optionally and each independently be
 replaced by halo, hydroxy, carboxyl, -P(=O)(OH)H, -P(=O)(OH)₂,
 -P(=O)(OH)CH₃, -P(=O)(OH)(OCH₃), -P(=O)(OH)(OCH₂CH₃),
 20 -P(=O)(OH)NH₂, -S(=O)₂H, -S(=O)₂(OH), -S(=O)₂NH,
 -C(=O)-NH-S(=O)₂-H, tetrazolyl, 3-hydroxy-isothiazolyl,
 3-hydroxy-isoxazolyl, 3-hydroxy-thiadiazolyl, mercaptotetrazolyl,
 3-mercapto-triazolyl, 3-sulfinyl-triazolyl, 3-sulfonyl-triazolyl;

or

25 L is a monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a
 bicyclic partially saturated or aromatic heterocycle wherein each of said ring
 systems may optionally be substituted with up to 3 substituents, each substituent
 independently being selected from C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy;
 aminocarbonyl; mono- or di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyl-C(=O)-NH-;
 30 C₁₋₆alkyloxy-C(=O)-NH-; H₂N-C(=O)-NH-; mono- or di(C₁₋₄alkyl)amino-C(=O)-
 NH-; Het-NH-; Het¹-NH-; -NH-C(=N-R^x)NR^yR^z; -C(=N-R^x)NR^yR^z; Het¹; or a
 radical of formula

-X-C₁-Y₁-C₂-Y₂-C₃-Y₃-C₄-Z (c-1) wherein

X represents NR⁵, O, S or a direct bond;

35 C₁, C₂, C₃ and C₄ each independently represent C₁₋₆alkanediyl,
 C₂₋₆alkenediyl, C₂₋₆alkynediyl or a direct bond;

Y₁, Y₂ and Y₃ each independently represent NR⁵, O, S or a direct bond;

Z is hydrogen, halo, cyano, hydroxy, carboxyl, -P(=O)(OH)H,
 -P(=O)(OH)₂, -P(=O)(OH)CH₃, -P(=O)(OH)(OCH₃),
 -P(=O)(OH)(OCH₂CH₃), -P(=O)(OH)NH₂, -S(=O)₂H, -S(=O)₂(OH),
 -S(=O)₂NH, -C(=O)-NH-S(=O)₂-H, tetrazolyl, 3-hydroxy-isothiazolyl,
 3-hydroxy-isoxazolyl, 3-hydroxy-thiadiazolyl, mercaptotetrazolyl,
 3-mercapto-triazolyl, 3-sulfinyl-triazolyl, 3-sulfonyl-triazolyl;
 R⁵ is hydrogen, C₁₋₆alkyl or -C(=NH)-N(R²)₂; and wherein
 from 1 to 3 hydrogen atoms of the C₁₋₆alkyl, C₁₋₆alkanediyl,
 C₂₋₆alkenediyl or C₂₋₆alkynediyl groups in the definitions of R⁵ and the
 radical of formula (c-1) may optionally and each independently be
 replaced by halo, hydroxy, carboxyl, -P(=O)(OH)H, -P(=O)(OH)₂,
 -P(=O)(OH)CH₃, -P(=O)(OH)(OCH₃), -P(=O)(OH)(OCH₂CH₃),
 -P(=O)(OH)NH₂, -S(=O)₂H, -S(=O)₂(OH), -S(=O)₂NH,
 -C(=O)-NH-S(=O)₂-H, tetrazolyl, 3-hydroxy-isothiazolyl,
 3-hydroxy-isoxazolyl, 3-hydroxy-thiadiazolyl; mercaptotetrazolyl,
 3-mercapto-triazolyl, 3-sulfinyl-triazolyl, 3-sulfonyl-triazolyl;


Het is a monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a
 bicyclic partially saturated or aromatic heterocycle wherein each of said ring
 systems may optionally be substituted with up to 3 substituents, each substituent
 independently being selected from halo, hydroxy, amino, cyano, carboxyl,
 mono-or di(C₁₋₆alkyl)amino, C₁₋₆alkyl, C₁₋₆alkyl substituted with hydroxy or
 C₁₋₄alkyloxy or amino or mono-or di(C₁₋₄alkyl)amino, polyhaloC₁₋₆alkyl,
 C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylcarbonyloxy,
 aminocarbonyl, mono-or di(C₁₋₆alkyl)aminocarbonyl, C₁₋₆alkyl-C(=O)-NH-,
 C₁₋₆alkyloxy-C(=O)-NH-, H₂N-C(=O)-NH- or mono- or di(C₁₋₄alkyl)amino-
 C(=O)-NH-;

Het¹ is a saturated 6-membered heterocycle selected from piperidinyl, morpholinyl,
 thiomorpholinyl, piperazinyl, wherein said saturated 6-membered heterocycle
 may optionally be substituted with amino or C₁₋₄alkyl optionally substituted
 with aryl;

aryl is phenyl, optionally substituted with up to five substituents each independently
 selected from halo, hydroxy, C₁₋₆alkyl, polyhaloC₁₋₆alkyl, C₁₋₆alkyloxy,
 C₁₋₆alkylthio, cyano, nitro, amino, mono-or di(C₁₋₆alkyl)amino.

The present invention also relates to a compound of formula


-6-



a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof,

wherein

- 5 Z is halo; C₁₋₆alkyl; C₁₋₆alkyl substituted with hydroxy, carboxyl, cyano, amino, mono -or di(C₁₋₆alkyl)amino, aminocarbonyl, mono -or di(C₁₋₆alkyl)aminocarbonyl, C₁₋₆alkyloxycarbonyl or C₁₋₆alkyloxy; polyhaloC₁₋₄alkyl; C₁₋₄alkyloxy; cyano; amino; aminocarbonyl; mono -or di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; H₂N-S(=O)₂-; mono -or
- 10 di(C₁₋₆alkyl)amino-S(=O)₂; -C(=N-R^x)NR^yR^z;
 R^x is hydrogen, C₁₋₆alkyl, cyano, nitro or -S(=O)₂-NH₂;
 R^y is hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl or C₂₋₆alkynyl;
 R^z is hydrogen or C₁₋₆alkyl;
- 15 Q is C₃₋₆cycloalkyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, phenyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl, or imidazopyridyl, wherein each of said ring systems may optionally be substituted with up to three substituents each of said
- 20 substituents independently being selected from halo; hydroxy; cyano; carboxyl; azido; amino; mono- or di(C₁₋₆alkyl)amino; C₁₋₆alkylcarbonylamino; C₁₋₆alkyl; C₂₋₆alkenyl; C₂₋₆alkynyl; C₃₋₆cycloalkyl; C₁₋₆alkyl substituted with hydroxy, C₁₋₆alkyloxy, amino, mono-or di(C₁₋₄alkyl)amino; C₁₋₆alkyloxy; C₁₋₆alkylthio; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; arylC₁₋₆alkyloxy; aryloxy; polyhaloC₁₋₆alkyl; polyhaloC₁₋₆alkyloxy; polyhaloC₁₋₆alkylcarbonyl;
- 25 C₁₋₄alkyl-S(=O)_n- or R¹HN-S(=O)_n-, with R¹ representing hydrogen, or a radical of formula




with A being O, S or a bivalent radical of formula -CR²=N- with CR² attached to N of formula (a-1); and

- 30 R² being hydrogen, C₁₋₆alkyl or C₁₋₆alkyloxy;

or

Q is a radical of formula

-7-



wherein B_1 and B_2 each independently are O, NR^3 , CH_2 or S, with R^3 being hydrogen or C_{1-4} alkyl;

B_3 is O or NR^4 with R^4 being hydrogen or C_{1-4} alkyl;

5 q is an integer with value 1 to 4;

r is an integer with value 1 to 3;

n is an integer with value 1 or 2;

10 L is a monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a bicyclic partially saturated or aromatic heterocycle wherein each of said ring systems may optionally be substituted with up to 3 substituents, each substituent independently being selected from C_{1-6} alkyloxycarbonyl; C_{1-6} alkylcarbonyloxy; aminocarbonyl; mono- or di(C_{1-6} alkyl)aminocarbonyl; C_{1-6} alkyl- $C(=O)$ -NH-; C_{1-6} alkyloxy- $C(=O)$ -NH-; H_2N - $C(=O)$ -NH-; mono- or di(C_{1-4} alkyl)amino- $C(=O)$ -NH-; Het-NH-; Het¹-NH-; -NH- $C(=N-R^x)NR^yR^z$; - $C(=N-R^x)NR^yR^z$; Het¹; or a radical of formula

15 $-X-C_1-Y_1-C_2-Y_2-C_3-Y_3-C_4-Z$ (c-1) wherein

X represents NR^5 , O, S or a direct bond;

C_1 , C_2 , C_3 and C_4 each independently represent C_{1-6} alkanediyl,

C_{2-6} alkenediyl, C_{2-6} alkynediyl or a direct bond;

20 Y_1 , Y_2 and Y_3 each independently represent NR^5 , O, S or a direct bond;

Z is hydrogen, halo, cyano, hydroxy, carboxyl, - $P(=O)(OH)H$,

- $P(=O)(OH)_2$, - $P(=O)(OH)CH_3$, - $P(=O)(OH)(OCH_3)$,

- $P(=O)(OH)(OCH_2CH_3)$, - $P(=O)(OH)NH_2$, - $S(=O)_2H$, - $S(=O)_2(OH)$,

25 - $S(=O)_2NH$, - $C(=O)$ -NH- $S(=O)_2$ -H, tetrazolyl, 3-hydroxy-isothiazolyl,

3-hydroxy-isoxazolyl, 3-hydroxy-thiadiazolyl, mercaptotetrazolyl,

3-mercapto-triazolyl, 3-sulfinyl-triazolyl, 3-sulfonyl-triazolyl;

R^5 is hydrogen, C_{1-6} alkyl or - $C(=NH)$ - $N(R^z)_2$; and wherein

from 1 to 3 hydrogen atoms of the C_{1-6} alkyl, C_{1-6} alkanediyl,

C_{2-6} alkenediyl or C_{2-6} alkynediyl groups in the definitions of R^5 and the

30 radical of formula (c-1) may optionally and each independently be replaced by halo, hydroxy, carboxyl, - $P(=O)(OH)H$, - $P(=O)(OH)_2$,

- $P(=O)(OH)CH_3$, - $P(=O)(OH)(OCH_3)$, - $P(=O)(OH)(OCH_2CH_3)$,

- $P(=O)(OH)NH_2$, - $S(=O)_2H$, - $S(=O)_2(OH)$, - $S(=O)_2NH$,

- $C(=O)$ -NH- $S(=O)_2$ -H, tetrazolyl, 3-hydroxy-isothiazolyl,

3-hydroxy-isoxazolyl, 3-hydroxy-thiadiazolyl, mercaptotetrazolyl,
3-mercapto-triazolyl, 3-sulfinyl-triazolyl, 3-sulfonyl-triazolyl;


Het is a monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a bicyclic partially saturated or aromatic heterocycle wherein each of said ring systems may optionally be substituted with up to 3 substituents, each substituent independently being selected from halo, hydroxy, amino, cyano, carboxyl, mono-or di(C₁₋₆alkyl)amino, C₁₋₆alkyl, C₁₋₆alkyl substituted with hydroxy or C₁₋₄alkyloxy or amino or mono-or di(C₁₋₄alkyl)amino, polyhaloC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylcarbonyloxy, aminocarbonyl, mono-or di(C₁₋₆alkyl)aminocarbonyl, C₁₋₆alkyl-C(=O)-NH-, C₁₋₆alkyloxy-C(=O)-NH-, H₂N-C(=O)-NH- or mono- or di(C₁₋₄alkyl)amino-C(=O)-NH-;















Het¹ is a saturated 6-membered heterocycle selected from piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, wherein said saturated 6-membered heterocycle may optionally be substituted with amino or C₁₋₄alkyl optionally substituted with aryl;

aryl is phenyl, optionally substituted with up to five substituents each independently selected from halo, hydroxy, C₁₋₆alkyl, polyhaloC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, cyano, nitro, amino, mono-or di(C₁₋₆alkyl)amino.


provided that

- when Z is methyl, Q is phenyl or phenyl substituted with halo, methyl or ethyloxy, then L is other than quinoxalinyll;
- when Z is methyl, Q is phenyl or phenyl substituted at the para position with methyl, chloro, nitro or methyloxy, then L is other than thiazolyl substituted with methyl or amino;
- when Z is trifluoromethyl, Q is 4-methylphenyl, then L is other than 1,2,3-triazolyl mono-or disubstituted with methyloxycarbonyl;
- L is other than unsubstituted or substituted benzoxazolyl or unsubstituted or substituted benzimidazolyl;
- the compound is other than

L	Z	Q
	CH ₃ -	phenyl
3-pyridyl	CH ₃ -	4-pyridyl

	$\text{CH}_3\text{-C(=O)-O-}$	phenyl
	$\text{CH}_3\text{-}$	4-chloro-phenyl
	$\text{CH}_3\text{-}$	phenyl
	-CF_3	4-methyl-phenyl
	$\text{-CH}_2\text{-C(=O)-O-CH}_2\text{CH}_3$	4-chloro-phenyl
	-CF_3	phenyl
	$\text{CH}_3\text{-}$	phenyl
	$\text{CH}_3\text{-}$	4-chlorophenyl
	$\text{CH}_3\text{-}$	phenyl
	$\text{CH}_3\text{-}$	2-thienyl
	F-	
	$\text{CH}_3\text{-}$	phenyl
	$\text{CH}_3\text{-}$	4-chlorophenyl

The present invention further relates to a compound of formula



a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a
 5 stereochemically isomeric form thereof,
 wherein


Z is halo; C₁₋₆alkyl; C₁₋₆alkyl substituted with hydroxy, carboxyl, cyano, amino, mono
 -or di(C₁₋₆alkyl)amino, aminocarbonyl, mono -or di(C₁₋₆alkyl)aminocarbonyl,
 C₁₋₆alkyloxycarbonyl or C₁₋₆alkyloxy; polyhaloC₁₋₄alkyl; C₁₋₄alkyloxy; cyano;
 10 amino; aminocarbonyl; mono -or di(C₁₋₆alkyl)aminocarbonyl;
 C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; H₂N-S(=O)₂-; mono -or
 di(C₁₋₆alkyl)amino-S(=O)₂; -C(=N-R^x)NR^yR^z;

R^x is hydrogen, C₁₋₆alkyl, cyano, nitro or -S(=O)₂-NH₂;

R^y is hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl or C₂₋₆alkynyl;

15 R^z is hydrogen or C₁₋₆alkyl;

Q is C₃₋₆cycloalkyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl,
 pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, phenyl,
 pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl,
 benzimidazolyl, indazolyl, or imidazopyridyl, wherein each of said ring systems
 20 may optionally be substituted with up to three substituents each of said
 substituents independently being selected from halo; hydroxy; cyano; carboxyl;
 azido; amino; mono- or di(C₁₋₆alkyl)amino; C₁₋₆alkylcarbonylamino; C₁₋₆alkyl;
 C₂₋₆alkenyl; C₂₋₆alkynyl; C₃₋₆cycloalkyl; C₁₋₆alkyl substituted with hydroxy,
 C₁₋₆alkyloxy, amino, mono-or di(C₁₋₄alkyl)amino; C₁₋₆alkyloxy; C₁₋₆alkylthio;
 25 C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; arylC₁₋₆alkyloxy; aryloxy;
 polyhaloC₁₋₆alkyl; polyhaloC₁₋₆alkyloxy; polyhaloC₁₋₆alkylcarbonyl;
 C₁₋₄alkyl-S(=O)_n- or R¹HN-S(=O)_n-, with R¹ representing hydrogen, or a radical
 of formula




30 with A being O, S or a bivalent radical of formula -CR²=N- with CR²
 attached to N of formula (a-1); and
 R² being hydrogen, C₁₋₆alkyl or C₁₋₆alkyloxy;

or

-11-

Q is a radical of formula



wherein B_1 and B_2 each independently are O, NR^3 , CH_2 or S, with R^3 being hydrogen or C_{1-4} alkyl;

B_3 is O or NR^4 with R^4 being hydrogen or C_{1-4} alkyl;

q is an integer with value 1 to 4;

r is an integer with value 1 to 3;

n is an integer with value 1 or 2;

L is a monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a

bicyclic partially saturated or aromatic heterocycle wherein each of said ring systems may optionally be substituted with up to 3 substituents, each substituent

independently being selected from C_{1-6} alkyloxycarbonyl; C_{1-6} alkylcarbonyloxy;

aminocarbonyl; mono- or di(C_{1-6} alkyl)aminocarbonyl; C_{1-6} alkyl-C(=O)-NH-;

C_{1-6} alkyloxy-C(=O)-NH-; $H_2N-C(=O)-NH-$; mono- or di(C_{1-4} alkyl)amino-C(=O)-

NH-; Het-NH-; Het¹-NH-; -NH-C(=N- R^x) NR^yR^z ; -C(=N- R^x) NR^yR^z ; Het¹; or a radical of formula

-X- C_1 - Y_1 - C_2 - Y_2 - C_3 - Y_3 - C_4 -Z (c-1) wherein

X represents NR^5 , O, S or a direct bond;

C_1 , C_2 , C_3 and C_4 each independently represent C_{1-6} alkanediyl,

C_{2-6} alkenediyl, C_{2-6} alkynediyl or a direct bond;

Y_1 , Y_2 and Y_3 each independently represent NR^5 , O, S or a direct bond;

Z is hydrogen, halo, cyano, hydroxy, carboxyl, -P(=O)(OH)H,

-P(=O)(OH)₂, -P(=O)(OH)CH₃, -P(=O)(OH)(OCH₃),

-P(=O)(OH)(OCH₂CH₃), -P(=O)(OH)NH₂, -S(=O)₂H, -S(=O)₂(OH),

-S(=O)₂NH, -C(=O)-NH-S(=O)₂-H, tetrazolyl, 3-hydroxy-isothiazolyl,

3-hydroxy-isoxazolyl, 3-hydroxy-thiadiazolyl, mercaptotetrazolyl,

3-mercapto-triazolyl, 3-sulfinyl-triazolyl, 3-sulfonyl-triazolyl;

R^5 is hydrogen, C_{1-6} alkyl or -C(=NH)-N(R^z)₂; and wherein

from 1 to 3 hydrogen atoms of the C_{1-6} alkyl, C_{1-6} alkanediyl,

C_{2-6} alkenediyl or C_{2-6} alkynediyl groups in the definitions of R^5 and the radical of formula (c-1) may optionally and each independently be

replaced by halo, hydroxy, carboxyl, -P(=O)(OH)H, -P(=O)(OH)₂,

-P(=O)(OH)CH₃, -P(=O)(OH)-(OCH₃), -P(=O)(OH)(OCH₂CH₃),

-P(=O)(OH)NH₂, -S(=O)₂H, -S(=O)₂(OH), -S(=O)₂NH,

-12-

-C(=O)-NH-S(=O)₂-H, tetrazolyl, 3-hydroxy-isothiazolyl,
 3-hydroxy-isoxazolyl, 3-hydroxy-thiadiazolyl, mercaptotetrazolyl,
 3-mercapto-triazolyl, 3-sulfinyl-triazolyl, 3-sulfonyl-triazolyl;


Het is a monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a
 5 bicyclic partially saturated or aromatic heterocycle wherein each of said ring
 systems may optionally be substituted with up to 3 substituents, each substituent
 independently being selected from halo, hydroxy, amino, cyano, carboxyl,
 mono-or di(C₁₋₆alkyl)amino, C₁₋₆alkyl, C₁₋₆alkyl substituted with hydroxy or
 C₁₋₄alkyloxy or amino or mono-or di(C₁₋₄alkyl)amino, polyhaloC₁₋₆alkyl,
 10 C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylcarbonyloxy,
 aminocarbonyl, mono-or di(C₁₋₆alkyl)aminocarbonyl, C₁₋₆alkyl-C(=O)-NH-,
 C₁₋₆alkyloxy-C(=O)-NH-, H₂N-C(=O)-NH- or mono- or di(C₁₋₄alkyl)amino-
 C(=O)-NH-;






Het¹ is a saturated 6-membered heterocycle selected from piperidinyl, morpholinyl,
 15 thiomorpholinyl, piperazinyl, wherein said saturated 6-membered heterocycle
 may optionally be substituted with amino or C₁₋₄alkyl optionally substituted
 with aryl;

aryl is phenyl, optionally substituted with up to five substituents each independently
 selected from halo, hydroxy, C₁₋₆alkyl, polyhaloC₁₋₆alkyl, C₁₋₆alkyloxy,
 20 C₁₋₆alkylthio, cyano, nitro, amino, mono-or di(C₁₋₆alkyl)amino.

provided that

- L is other than substituted phenyl;
- when Z is methyl, Q is phenyl or phenyl substituted with halo, CH₃ or ethyloxy, then L
 is other than quinoxalinyl;
- 25 -when Z is methyl, Q is phenyl or phenyl substituted at the para position with methyl,
 chloro, nitro or methyloxy, then L is other than thiazolyl substituted with methyl or
 amino;
- the compound is other than

L	Z	Q
3-pyridyl	CH ₃ -	4-pyridyl
	CH ₃ -	4-chloro-phenyl

	CH ₃ -	phenyl
	-CH ₂ -C(=O)-O-CH ₂ CH ₃	4-chloro-phenyl
	CH ₃ -	phenyl
	F-	

for use as a medicine.

- As used hereinabove or hereinafter C₁₋₄alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as methyl, ethyl, propyl, 1-methylethyl, butyl, 2-methylpropyl and the like; C₁₋₆alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as the groups defined for C₁₋₄alkyl and pentyl, hexyl, 2-methylbutyl, 3-methylpentyl and the like; C₁₋₁₂alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 12 carbon atoms such as the groups defined for C₁₋₆alkyl and heptyl, octyl, nonyl, decyl, 3-ethylpentyl and the like; C₁₋₆alkanediyl defines straight or branched chain saturated bivalent hydrocarbon radicals having from 1 to 6 carbon atoms such as methylene, 1,2-ethanediyl or 1,2-ethylidene, 1,3-propanediyl or 1,3-propylidene, 1,4-butanediyl or 1,4-butylidene, 1,5-pentanediyl, 1,6-hexanediyl and the like; C₂₋₆alkenyl as a group or part of a group defines straight or branched chain hydrocarbon radicals having from 2 to 6 carbon atoms and having 1 double bond such as ethenyl, propenyl, butenyl, pentenyl, hexenyl, 3-methylbutenyl and the like; C₂₋₆alkenediyl as a group or part of a group defines straight or branched chain bivalent hydrocarbon radicals having from 2 to 6 carbon atoms and having 1 double bond such as ethenediyl, 2-butene-1,4-diyl and the like; C₂₋₆alkynyl as a group or part of a group defines straight or branched chain hydrocarbon radicals having from 2 to 6 carbon atoms and having 1 triple bond such as ethynyl, propynyl, butynyl, pentynyl, hexynyl, 3-methylbutynyl and the like; C₂₋₆alkynediyl as a group or part of a group defines straight or branched chain bivalent hydrocarbon radicals having from 2 to 6 carbon atoms and having 1 triple bond such as ethynediyl, 3-pentyne-1,5-diyl and the like; C₃₋₆cycloalkyl is generic to cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl; a

monocyclic or bicyclic partially saturated heterocycle represents a ring system consisting of 1 or 2 rings and comprising at least one heteroatom selected from O, N or S, and at least one double bond provided that the ring system is not an aromatic system; a monocyclic or bicyclic aromatic heterocycle represents an aromatic ring system
5 consisting of 1 or 2 rings and comprising at least one heteroatom selected from O, N or S; the term aromatic is well known to a person skilled in the art and designates cyclically conjugated systems of $4n' + 2$ electrons, that is with 6, 10, 14 etc. π -electrons (rule of Hückel).

- 10 The L or Q radical as described above for the compounds of formula (I) may be attached to the remainder of the molecule of formula (I) through any ring carbon or heteroatom as appropriate. For example, when Q is pyridyl, it may be 2-pyridyl, 3-pyridyl or 4-pyridyl.
- 15 Lines drawn into ring systems indicate that the bond may be attached to any suitable ring atom. When the ring system is a bicyclic ring system, the bond may be attached to any suitable ring atom of either of the two rings.

- As used herein before, the term (=O) forms a carbonyl moiety when attached to a
20 carbon atom, a sulfoxide moiety when attached to a sulfur atom and a sulfonyl moiety when two of said terms are attached to a sulfur atom.

- The term halo is generic to fluoro, chloro, bromo and iodo. As used in the foregoing and hereinafter, polyhaloC₁₋₆alkyl as a group or part of a group is defined as mono- or
25 polyhalosubstituted C₁₋₆alkyl, in particular methyl with one or more fluoro atoms, for example, difluoromethyl or trifluoromethyl. In case more than one halogen atoms are attached to an alkyl group within the definition of polyhaloC₁₋₆alkyl, they may be the same or different.

- 30 When any variable occurs more than one time in any constituent, each definition is independent.

- Whenever used hereinbefore or hereinafter that substituents can be selected each independently out of a list of numerous definitions, such as for example the
35 substituents of L or Q, all possible combinations are intended which are chemically possible and which lead to chemically stable molecules. When ring systems are attached to the remainder of the molecule via a linker, such as for example Het¹-NH-,

all possible combinations of ring system and linker are intended which are chemically possible and which lead to chemically stable molecules.

When several consecutive substituents in the radical of formula (c-1) represent a direct bond, this has to be interpreted as one single direct bond. For instance, when X represents a direct bond, C₁ represents CH₂, Y₁, C₂, Y₂, C₃, Y₃ and C₄ represent a direct bond and Z represents hydrogen, then said radical of formula (c-1) represents methyl (CH₃).

It will be appreciated that some of the compounds of formula (I) and their *N*-oxides, addition salts, quaternary amines and stereochemically isomeric forms may contain one or more centers of chirality and exist as stereochemically isomeric forms.

The term "stereochemically isomeric forms" as used hereinbefore or hereinafter defines all the possible stereoisomeric forms which the compounds of formula (I) and their *N*-oxides, addition salts, quaternary amines or physiologically functional derivatives may possess. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture of all possible stereochemically isomeric forms, said mixtures containing all diastereomers and enantiomers of the basic molecular structure as well as each of the individual isomeric forms of formula (I) and their *N*-oxides, salts, solvates, quaternary amines substantially free, *i.e.* associated with less than 10%, preferably less than 5%, in particular less than 2% and most preferably less than 1% of the other isomers. Stereochemically isomeric forms of the compounds of formula (I) are obviously intended to be embraced within the scope of this invention.

For therapeutic use, salts of the compounds of formula (I) are those wherein the counterion is pharmaceutically acceptable. However, salts of acids and bases which are non-pharmaceutically acceptable may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound. All salts, whether pharmaceutically acceptable or not are included within the ambit of the present invention.

The pharmaceutically acceptable acid and base addition salts as mentioned hereinabove or hereinafter are meant to comprise the therapeutically active non-toxic acid and base addition salt forms which the compounds of formula (I) are able to form. The pharmaceutically acceptable acid addition salts can conveniently be obtained by treating the base form with such appropriate acid. Appropriate acids comprise, for example,

inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid, sulfuric, nitric, phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic (i.e. ethanedioic), malonic, succinic (i.e. butanedioic acid), maleic, fumaric, malic, tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-aminosalicylic, pamoic and the like acids.

Conversely said salt forms can be converted by treatment with an appropriate base into the free base form.

- 10 The compounds of formula (I) containing an acidic proton may also be converted into their non-toxic metal or amine addition salt forms by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g.
- 15 primary, secondary and tertiary aliphatic and aromatic amines such as methylamine, ethylamine, propylamine, isopropylamine, the four butylamine isomers, dimethylamine, diethylamine, diethanolamine, dipropylamine, diisopropylamine, di-*n*-butylamine, pyrrolidine, piperidine, morpholine, trimethylamine, triethylamine, tripropylamine, quinuclidine, pyridine, quinoline and isoquinoline; the benzathine, *N*-methyl-D-glucamine, hydrabamine salts, and salts with amino acids such as, for
- 20 example, arginine, lysine and the like.

Conversely the salt form can be converted by treatment with acid into the free acid form.

- 25 The term addition salt as used hereinabove also comprises the solvates which the compounds of formula (I) as well as the salts thereof, are able to form. Such solvates are for example hydrates, alcoholates and the like.

- The term "quaternary amine" as used hereinbefore defines the quaternary ammonium salts which the compounds of formula (I) are able to form by reaction between a basic nitrogen of a compound of formula (I) and an appropriate quaternizing agent, such as, for example, an optionally substituted alkylhalide, arylhalide or arylalkylhalide, e.g. methyl iodide or benzyl iodide. Other reactants with good leaving groups may also be used, such as alkyl trifluoromethanesulfonates, alkyl methanesulfonates, and alkyl
- 30 *p*-toluenesulfonates. A quaternary amine has a positively charged nitrogen.

Pharmaceutically acceptable counterions include for example chloro, bromo, iodo,

trifluoroacetate and acetate. The counterion of choice can be made using ion exchange resin columns.

5 The *N*-oxide forms of the present compounds are meant to comprise the compounds of formula (I) wherein one or several tertiary nitrogen atoms are oxidized to the so-called *N*-oxide.

10 Some of the compounds of formula (I) may also exist in their tautomeric form. Such forms although not explicitly indicated in the above formula are intended to be included within the scope of the present invention.


Particular examples of monocyclic or bicyclic partially saturated heterocycles are pyrrolinyl, imidazolinyl, pyrazolinyl, 2,3-dihydrobenzofuranyl, 1,3-benzodioxolyl, 2,3-dihydro-1,4-benzodioxinyl, indolinyl and the like.

15 Particular examples of monocyclic or bicyclic aromatic heterocycles are azetyl, oxetylidenyl, pyrrolyl, furyl, thienyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, triazolyl, thiadiazolyl, oxadiazolyl, tetrazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, pyranyl, benzofuryl, isobenzofuryl, 20 benzothienyl, isobenzothienyl, indoliziny, indolyl, isoindolyl, benzoxazolyl, benzimidazolyl, indazolyl, benzisoxazolyl, benzisothiazolyl, benzopyrazolyl, benzoxadiazolyl, benzothiadiazolyl, benzotriazolyl, purinyl, quinolinyl, isoquinolinyl, cinnolinyl, quinoliziny, phthalazinyl, quinoxaliny, quinazolinyl, naphthiridinyl, pteridinyl, benzopyranyl, pyrrolopyridyl, thienopyridyl, furopyridyl, isothiazolopyridyl, 25 thiazolopyridyl, isoxazolopyridyl, oxazolopyridyl, pyrazolopyridyl, imidazopyridyl, pyrrolopyrazinyl, thienopyrazinyl, furopyrazinyl, isothiazolopyrazinyl, thiazolopyrazinyl, isoxazolopyrazinyl, oxazolopyrazinyl, pyrazolopyrazinyl, imidazopyrazinyl, pyrrolopyrimidinyl, thienopyrimidinyl, fuopyrimidinyl, isothiazolopyrimidinyl, thiazolopyrimidinyl, isoxazolopyrimidinyl, 30 oxazolopyrimidinyl, pyrazolopyrimidinyl, imidazopyrimidinyl, pyrrolopyridazinyl, thienopyridazinyl, fuopyridazinyl, isothiazolopyridazinyl, thiazolopyridazinyl, isoxazolopyridazinyl, oxazolopyridazinyl, pyrazolopyridazinyl, imidazopyridazinyl, oxadiazolopyridyl, thiadiazolopyridyl, triazolopyridyl, oxadiazolopyrazinyl, thiadiazolopyrazinyl, triazolopyrazinyl, oxadiazolopyrimidinyl, thiadiazolopyrimidinyl, 35 triazolopyrimidinyl, oxadiazolopyridazinyl, thiadiazolopyridazinyl, triazolopyridazinyl, imidazooxazolyl, imidazothiazolyl, imidazoimidazolyl, isoxazolotriazinyl, isothiazolo-

triazinyl, pyrazolotriazinyl, oxazolotriazinyl, thiazolotriazinyl, imidazotriazinyl, oxadiazolotriazinyl, thiadiazolotriazinyl, triazolotriazinyl.

An interesting embodiment of the present invention concerns the use of those
 5 compounds of formula (I) wherein Q is C₃₋₆cycloalkyl, phenyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl, or imidazopyridyl, each of said rings optionally being substituted with up to three substituents each independently selected from halo; hydroxy; cyano; carboxyl; azido; amino; mono- or di(C₁₋₆alkyl)amino; C₁₋₆alkylcarbonylamino; C₁₋₆alkyl; C₂₋₆alkenyl; C₂₋₆alkynyl; C₃₋₆cycloalkyl; C₁₋₆alkyl substituted with hydroxy, C₁₋₆alkyloxy, amino, mono-or di(C₁₋₄alkyl)amino; C₁₋₆alkyloxy; C₁₋₆alkylthio; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; arylC₁₋₆alkyloxy; aryloxy; polyhaloC₁₋₆alkyl; polyhalo-C₁₋₆alkyloxy; polyhaloC₁₋₆alkylcarbonyl; C₁₋₄alkyl-S(=O)_n- or R¹HN-S(=O)_n-;
 10 or

15 Q is a radical of formula



wherein X and Y each independently are O, NR³, CH₂ or S, with R³ being hydrogen or C₁₋₄alkyl;

q is an integer with value 1 to 4;

20 Z is O or NR⁴ with R⁴ being hydrogen or C₁₋₄alkyl;

r is an integer with value 1 to 3;


Z is halo; C₁₋₆alkyl; C₁₋₆alkyl substituted with hydroxy, carboxyl, cyano, amino, mono-or di(C₁₋₆alkyl)amino, aminocarbonyl, mono-or di(C₁₋₆alkyl)aminocarbonyl, C₁₋₆alkyloxycarbonyl, C₁₋₆alkyloxy; polyhaloC₁₋₄alkyl; cyano; amino; aminocarbonyl; mono-or di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; aminoS(=O)₂-; mono-or di(C₁₋₆alkyl)aminoS(=O)₂; -C(=N-R^x)NR^yR^z;

L is phenyl, substituted with up to 4 substituents each independently being selected from halo, hydroxy, mercapto, amino, cyano, carboxyl, mono-or di(C₁₋₆alkyl)amino, C₁₋₆alkyl, C₁₋₆alkyl substituted with hydroxy or C₁₋₄alkyloxy or amino or mono-or di(C₁₋₄alkyl)amino, polyhaloC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylcarbonyloxy, aminocarbonyl, mono-or di(C₁₋₆alkyl)aminocarbonyl, C₁₋₆alkyl-C(=O)-NH-, C₁₋₆alkyloxy-C(=O)-NH-, H₂N-C(=O)-NH-, mono- or di(C₁₋₄alkyl)amino-C(=O)-NH- or Het-NH-, -C(=N-R^x)NR^yR^z; or

L is a monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a bicyclic partially saturated or aromatic heterocycle wherein each of said ring systems may optionally be substituted with up to 3 substituents, each substituent independently being selected from halo, hydroxy, mercapto, amino, cyano, carboxyl, mono-or di(C₁₋₆alkyl)amino, C₁₋₆alkyl, C₁₋₆alkyl substituted with hydroxy or C₁₋₄alkyloxy or amino or mono-or di(C₁₋₄alkyl)amino, polyhaloC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylcarbonyloxy, aminocarbonyl, mono-or di(C₁₋₆alkyl)aminocarbonyl, C₁₋₆alkyl-C(=O)-NH-, C₁₋₆alkyloxy-C(=O)-NH-, H₂N-C(=O)-NH-, mono- or di(C₁₋₄alkyl)amino-C(=O)-NH- or Het-NH-, -C(=N-R^x)NR^yR^z.

Another interesting embodiment of the present invention concerns those compounds of formula (I) wherein Q is C₃₋₆cycloalkyl, phenyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl, or imidazopyridyl, each of said rings optionally being substituted with up to three substituents each independently selected from halo; hydroxy; cyano; carboxyl; azido; amino; mono- or di(C₁₋₆alkyl)amino; C₁₋₆alkylcarbonylamino; C₁₋₆alkyl; C₂₋₆alkenyl; C₂₋₆alkynyl; C₃₋₆cycloalkyl; C₁₋₆alkyl substituted with hydroxy, C₁₋₆alkyloxy, amino, mono-or di(C₁₋₄alkyl)amino; C₁₋₆alkyloxy; C₁₋₆alkylthio; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; arylC₁₋₆alkyloxy; aryloxy; polyhaloC₁₋₆alkyl; polyhalo-C₁₋₆alkyloxy; polyhaloC₁₋₆alkylcarbonyl; C₁₋₄alkyl-S(=O)_n- or R¹HN-S(=O)_n-; or

Q is a radical of formula



wherein X and Y each independently are O, NR³, CH₂ or S, with R³ being hydrogen or C₁₋₄alkyl;
 q is an integer with value 1 to 4;
 Z is O or NR⁴ with R⁴ being hydrogen or C₁₋₄alkyl;
 r is an integer with value 1 to 3;
 Z is halo; C₁₋₆alkyl; C₁₋₆alkyl substituted with hydroxy, carboxyl, cyano, amino, mono-or di(C₁₋₆alkyl)amino, aminocarbonyl, mono-or di(C₁₋₆alkyl)aminocarbonyl, C₁₋₆alkyloxycarbonyl, C₁₋₆alkyloxy; polyhaloC₁₋₄alkyl; cyano; amino; aminocarbonyl; mono-or di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; aminoS(=O)₂-; mono-or di(C₁₋₆alkyl)aminoS(=O)₂; -C(=N-R^x)NR^yR^z;

L is a monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a bicyclic partially saturated or aromatic heterocycle wherein each of said ring systems may optionally be substituted with up to 3 substituents, each substituent independently being selected from halo, hydroxy, mercapto, amino, cyano, carboxyl, mono-or di(C₁₋₆alkyl)amino, C₁₋₆alkyl, C₁₋₆alkyl substituted with hydroxy or C₁₋₄alkyloxy or amino or mono-or di(C₁₋₄alkyl)amino, polyhaloC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylcarbonyloxy, aminocarbonyl, mono-or di(C₁₋₆alkyl)aminocarbonyl, C₁₋₆alkyl-C(=O)-NH-, C₁₋₆alkyloxy-C(=O)-NH-, H₂N-C(=O)-NH-, mono- or di(C₁₋₄alkyl)amino-C(=O)-NH- or Het-NH-, -C(=N-R^x)NR^yR^z.

A further interesting embodiment of the present invention concerns the use of those compounds of formula (I) wherein L is a monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a bicyclic partially saturated or aromatic heterocycle wherein each of said ring systems may optionally be substituted with up to 3 substituents, each substituent independently being selected from C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; aminocarbonyl; mono-or di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyl-C(=O)-NH-; C₁₋₆alkyloxy-C(=O)-NH-; H₂N-C(=O)-NH-; mono- or di(C₁₋₄alkyl)amino-C(=O)-NH-; Het-NH-; Het¹-NH-; -NH-C(=N-R^x)NR^yR^z; -C(=N-R^x)NR^yR^z; Het¹; or a radical of formula -X-C₁-Y₁-C₂-Y₂-C₃-Y₃-C₄-Z (c-1).

Also an interesting embodiment of the present invention concerns those compounds of formula (I) wherein one or, wherever possible, more of the following restrictions apply :

a) Q is C₃₋₆cycloalkyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, phenyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl, or imidazopyridyl, wherein each of said ring systems may optionally be substituted with up to three substituents each of said substituents independently being selected from halo; hydroxy; cyano; carboxyl; azido; amino; mono- or di(C₁₋₆alkyl)-amino; C₁₋₆alkylcarbonylamino; C₁₋₆alkyl; C₂₋₆alkenyl; C₂₋₆alkynyl; C₃₋₆cycloalkyl; C₁₋₆alkyl substituted with hydroxy, C₁₋₆alkyloxy, amino, mono-or di(C₁₋₄alkyl)amino; C₁₋₆alkyloxy; C₁₋₆alkylthio; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; arylC₁₋₆alkyloxy; aryloxy; polyhaloC₁₋₆alkyl; polyhaloC₁₋₆alkyloxy; polyhaloC₁₋₆alkylcarbonyl or C₁₋₄alkyl-S(=O)_n-; or Q is pyridyl substituted with up to three substituents each independently selected from halo; hydroxy; cyano; carboxyl; azido; amino; mono- or

- di(C₁₋₆alkyl)amino; C₁₋₆alkylcarbonylamino; C₁₋₆alkyl; C₂₋₆alkenyl; C₂₋₆alkynyl; C₃₋₆cycloalkyl; C₁₋₆alkyl substituted with hydroxy, C₁₋₆alkyloxy, amino, mono-or di(C₁₋₄alkyl)amino; C₁₋₆alkyloxy; C₁₋₆alkylthio; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; arylC₁₋₆alkyloxy; aryloxy; polyhaloC₁₋₆alkyl; polyhaloC₁₋₆alkyloxy; polyhaloC₁₋₆alkylcarbonyl or C₁₋₄alkyl-S(=O)_n-; b) Q is C₃₋₆cycloalkyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, phenyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl, or imidazopyridyl, wherein each of said ring systems may optionally be substituted with up to three substituents each of said substituents independently being selected from halo; hydroxy; cyano; carboxyl; azido; amino; mono- or di(C₁₋₆alkyl)amino; C₁₋₆alkylcarbonylamino; C₁₋₆alkyl; C₂₋₆alkenyl; C₂₋₆alkynyl; C₃₋₆cycloalkyl; C₁₋₆alkyl substituted with hydroxy, C₁₋₆alkyloxy, amino, mono-or di(C₁₋₄alkyl)amino; C₁₋₆alkyloxy; C₁₋₆alkylthio; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; arylC₁₋₆alkyloxy; aryloxy; polyhaloC₁₋₆alkyl; polyhaloC₁₋₆alkyloxy; polyhaloC₁₋₆alkylcarbonyl; C₁₋₄alkyl-S(=O)_n-; c) Q is furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, phenyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl, or imidazopyridyl, wherein each of said ring systems may optionally be substituted with up to three substituents each of said substituents independently being selected from halo; hydroxy; cyano; carboxyl; azido; amino; mono- or di(C₁₋₆alkyl)amino; C₁₋₆alkylcarbonylamino; C₁₋₆alkyl; C₂₋₆alkenyl; C₂₋₆alkynyl; C₃₋₆cycloalkyl; C₁₋₆alkyl substituted with hydroxy, C₁₋₆alkyloxy, amino, mono-or di(C₁₋₄alkyl)amino; C₁₋₆alkyloxy; C₁₋₆alkylthio; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; arylC₁₋₆alkyloxy; aryloxy; polyhaloC₁₋₆alkyl; polyhaloC₁₋₆alkyloxy; polyhaloC₁₋₆alkylcarbonyl; C₁₋₄alkyl-S(=O)_n-; d) Q is C₃₋₆cycloalkyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, phenyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl, or imidazopyridyl, wherein each of said ring systems may optionally be substituted with up to three substituents each of said substituents independently being selected from halo; cyano; carboxyl; azido; amino; mono- or di(C₁₋₆alkyl)amino; C₁₋₆alkylcarbonylamino; C₁₋₆alkyl; C₂₋₆alkenyl; C₂₋₆alkynyl; C₃₋₆cycloalkyl; C₁₋₆alkyl substituted with hydroxy, C₁₋₆alkyloxy, amino, mono-or di(C₁₋₄alkyl)amino; C₁₋₆alkyloxy; C₁₋₆alkylthio; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; arylC₁₋₆alkyloxy;

- aryloxy; polyhaloC₁₋₆alkyl; polyhaloC₁₋₆alkyloxy; polyhaloC₁₋₆alkylcarbonyl;
C₁₋₄alkyl-S(=O)_n-;
- e) Q is C₃₋₆cycloalkyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, phenyl, pyridyl,
5 pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl, or imidazopyridyl, wherein each of said ring systems may optionally be substituted with up to three substituents each of said substituents independently being selected from halo; hydroxy; cyano; carboxyl; azido; amino; mono- or di(C₁₋₆alkyl)-amino; C₁₋₆alkylcarbonylamino; C₁₋₆alkyl; C₂₋₆alkenyl; C₂₋₆alkynyl;
10 C₃₋₆cycloalkyl; C₁₋₆alkyl substituted with hydroxy, C₁₋₆alkyloxy, amino, mono-or di(C₁₋₄alkyl)amino; C₁₋₆alkyloxy; C₁₋₆alkylthio; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; arylC₁₋₆alkyloxy; aryloxy; polyhaloC₁₋₆alkyl; polyhaloC₁₋₆alkyloxy; polyhaloC₁₋₆alkylcarbonyl; C₁₋₄alkyl-S(=O)_n- or R¹HN-S(=O)_n-;
- f) L is furanyl, pyrrolyl, oxazolyl, imidazolyl, pyrazolyl, isothiazolyl, oxadiazolyl,
15 triazolyl, thiadiazolyl, a 5-membered partially saturated heterocycle, a 6-membered partially saturated or aromatic heterocycle or a bicyclic partially saturated or aromatic heterocycle wherein each of the aforesaid ring systems may optionally be substituted with up to 3 substituents, each substituent independently being selected from C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; aminocarbonyl; mono-or
20 di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyl-C(=O)-NH-; C₁₋₆alkyloxy-C(=O)-NH-; H₂N-C(=O)-NH-; mono- or di(C₁₋₄alkyl)amino-C(=O)-NH-; Het-NH-; Het¹-NH-; -NH-C(=N-R^x)NR^yR^z; -C(=N-R^x)NR^yR^z; Het¹; or a radical of formula -X-C₁-Y₁-C₂-Y₂-C₃-Y₃-C₄-Z (c-1); provided L is other than optionally substituted quinoxaliny;
- g) L is a 6-membered partially saturated or aromatic heterocycle wherein each of said ring systems may optionally be substituted with up to 3 substituents, each substituent independently being selected from C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; aminocarbonyl; mono-or di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyl-C(=O)-NH-;
C₁₋₆alkyloxy-C(=O)-NH-; H₂N-C(=O)-NH-; mono- or di(C₁₋₄alkyl)amino-C(=O)-NH-;
30 Het-NH-; Het¹-NH-; -NH-C(=N-R^x)NR^yR^z; -C(=N-R^x)NR^yR^z; Het¹; or a radical of formula -X-C₁-Y₁-C₂-Y₂-C₃-Y₃-C₄-Z (c-1);
- h) L is a monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a bicyclic partially saturated or aromatic heterocycle wherein each of said ring systems may optionally be substituted with up to 3 substituents, each substituent independently
35 being selected from C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; aminocarbonyl; mono-or di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyl-C(=O)-NH-; C₁₋₆alkyloxy-C(=O)-NH-;

- H₂N-C(=O)-NH-; mono- or di(C₁₋₄alkyl)amino-C(=O)-NH-; -NH-C(=N-R^x)NR^yR^z; -C(=N-R^x)NR^yR^z; Het¹; or a radical of formula -X-C₁-Y₁-C₂-Y₂-C₃-Y₃-C₄-Z (c-1);
- i) L is a monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a bicyclic partially saturated or aromatic heterocycle wherein each of said ring systems may optionally be substituted with up to 3 substituents, each substituent independently being selected from a radical of formula -X-C₁-Y₁-C₂-Y₂-C₃-Y₃-C₄-Z (c-1);
- j) L is a monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a bicyclic partially saturated or aromatic heterocycle wherein each of said ring systems may optionally be substituted with up to 3 substituents, each substituent independently being selected from a radical of formula -X-C₁-Y₁-C₂-Y₂-C₃-Y₃-C₄-Z (c-1) provided said radical does not represent hydroxy;
- k) L is a monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a bicyclic partially saturated or aromatic heterocycle wherein each of said ring systems may optionally be substituted with 1 or 2 substituents, each of said two substituent independently being selected from C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; aminocarbonyl; mono-or di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyl-C(=O)-NH-; C₁₋₆alkyloxy-C(=O)-NH-; H₂N-C(=O)-NH-; mono- or di(C₁₋₄alkyl)amino-C(=O)-NH-; Het-NH-; Het¹-NH-; -NH-C(=N-R^x)NR^yR^z; -C(=N-R^x)NR^yR^z; Het¹; or a radical of formula -X-C₁-Y₁-C₂-Y₂-C₃-Y₃-C₄-Z (c-1);
- l) L is a monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a bicyclic partially saturated or aromatic heterocycle wherein each of said ring systems may optionally be substituted with up to 3 substituents, each substituent independently being selected from halo; hydroxy; mercapto; amino; cyano; carboxyl; mono-or di(C₁₋₁₂alkyl)amino optionally being substituted with one, two or three substituents each independently being selected from hydroxy, amino, mono-or di(C₁₋₄alkyl)amino, C₁₋₄alkyloxy, C₁₋₄alkyloxyC₁₋₄alkyloxy, aminoC₁₋₄alkyloxy, aminoC₁₋₄alkyloxyC₁₋₄alkyloxy, mono-or di(C₁₋₄alkyl)aminoC₁₋₄alkyloxy, mono-or di(C₁₋₄alkyl)aminoC₁₋₄alkyloxyC₁₋₄alkyloxy; C₁₋₆alkyl; C₁₋₆alkyl substituted with hydroxy, C₁₋₄alkyloxy, amino or mono-or di(C₁₋₄alkyl)amino; polyhaloC₁₋₆alkyl; C₁₋₆alkyloxy; C₁₋₆alkylthio; C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; aminocarbonyl; mono-or di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyl-C(=O)-NH-; C₁₋₆alkyloxy-C(=O)-NH-; H₂N-C(=O)-NH-; mono- or di(C₁₋₄alkyl)amino-C(=O)-NH-; Het-NH-; Het¹-NH-; -NH-C(=N-R^x)NR^yR^z; -C(=N-R^x)NR^yR^z or Het¹;
- m) Z is halo; C₁₋₆alkyl; C₁₋₆alkyl substituted with hydroxy, carboxyl, cyano, amino, mono-or di(C₁₋₆alkyl)amino, aminocarbonyl, mono-or di(C₁₋₆alkyl)aminocarbonyl, C₁₋₆alkyloxycarbonyl or C₁₋₆alkyloxy; polyhaloC₁₋₄alkyl; cyano; aminocarbonyl; mono

-or di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy;
H₂N-S(=O)₂-; mono -or di(C₁₋₆alkyl)aminoS(=O)₂; -C(=N-R^x)NR^yR^z.

Also an interesting embodiment of the present invention concerns those compounds of
5 formula (I) wherein L is a substituted aromatic 6-membered heterocycle, in particular
substituted pyrimidinyl or substituted triazinyl, more in particular substituted
pyrimidin-4-yl, even more in particular pyrimidin-4-yl substituted with 1 or 2
substituents, said substituents preferably being selected from amino,
aminoC₁₋₆alkylamino, hydroxyC₁₋₆alkylamino (e.g. 3-hydroxypropylamino),
10 carboxyC₁₋₆alkylamino (e.g. 2-carboxyethylamino) or halo (e.g. fluoro).

A further interesting embodiment of the present invention concerns those compounds
of formula (I) wherein Q is phenyl; phenyl substituted with one or two substituents
selected from halo, polyhaloC₁₋₆alkyl; pyridyl; Z is C₁₋₆alkyl or halo; L is pyrimidinyl,
15 pyrazolyl or triazolyl, each of said three rings being optionally substituted with one or
two substituents selected from halo, amino, C₁₋₆alkylcarbonylamino,
C₁₋₆alkylamino, C₁₋₆alkylthio, Het-NH-.

Also a preferred embodiment of the present invention concerns those compounds of
20 formula (I) wherein Q is phenyl substituted with polyhaloC₁₋₆alkyl, in particular
trifluoromethyl.

Also preferred are those compounds of formula (I) wherein Z is halo or C₁₋₆alkyl, in
particular chloro or methyl.

Also a preferred embodiment of the present invention concerns those compounds of
formula (I) wherein Q is phenyl, pyridyl, pyrrolyl, pyrazolyl or thienyl, wherein each of
said ring systems may optionally be substituted with one or two substituents each
independently being selected from halo or polyhaloC₁₋₆alkyl; Z is C₁₋₆alkyl, halo,
30 C₁₋₆alkyloxy, aminocarbonyl; L is pyrimidinyl, pyrazolyl, triazolyl or triazinyl, wherein
each of said ring systems may optionally be substituted with one or two substituents
each independently being selected from amino, C₁₋₆alkylcarbonylamino, halo, Het-NH-
, hydroxy, C₁₋₆alkylthio, C₁₋₆alkyloxy, C₁₋₆alkyl, C₁₋₁₂alkylamino, mono-or
di(hydroxyC₁₋₁₂alkyl)amino wherein C₁₋₁₂alkyl may further optionally be substituted
35 with hydroxy, Het¹, aminocarbonyl, cyano, aminoC₁₋₁₂alkylamino,
hydroxyC₁₋₁₂alkyloxy, -NH-C(=NH)-NH₂, carboxyC₁₋₁₂alkylamino or
aminoC₁₋₆alkyloxyC₁₋₆alkyloxyC₁₋₆alkylamino.

Still a further interesting embodiment of the present invention concerns those compounds of formula (I) wherein Q is phenyl; phenyl substituted with one or two substituents selected from chloro, fluoro, trifluoromethyl; 2-pyridyl or 3-pyridyl; Z is methyl; L is 2-amino-pyrimidin-4-yl, 2-methylcarbonylamino-pyrimidin-4-yl, 2-methylamino-pyrimidin-4-yl, 2-methylthio-pyrimidin-4-yl, 2-[(4-methyl-2-thiazolyl)amino]-pyrimidin-4-yl, 2-amino-5-bromo-pyrimidin-4-yl, 2-amino-5-chloro-pyrimidin-4-yl, 4-pyrimidinyl, 3-pyrazolyl, 2-methylthio-1-methyl-1,3,4-triazol-5-yl.

10

Another interesting embodiment of the present invention concerns those compounds of formula (I) wherein Q is pyrrolyl; pyrazolyl; thienyl; phenyl; phenyl substituted with one or two substituents selected from chloro, fluoro, methyl, trifluoromethyl; 2-pyridyl; 2-pyridyl substituted with chloro, fluoro or trifluoromethyl; 3-pyridyl; Z is methyl,

15

chloro, methoxy or aminocarbonyl; L is 2-amino-pyrimidin-4-yl, 2-methylcarbonylamino-pyrimidin-4-yl, 2-methylamino-pyrimidin-4-yl, 2-methylthio-pyrimidin-4-yl, 2-[(4-methyl-2-thiazolyl)amino]-pyrimidin-4-yl, 2-amino-5-bromo-pyrimidin-4-yl, 2-amino-5-chloro-pyrimidin-4-yl, 2-amino-5-fluoro-pyrimidin-4-yl, 2-(2-hydroxy)ethylamino-pyrimidin-4-yl, 2-(3-hydroxy)propylamino-pyrimidin-4-yl, 2-piperazinyl-pyrimidin-4-yl, 2-(4-methylpiperazinyl)-pyrimidin-4-yl, 2-(2-amino)ethylamino-pyrimidin-4-yl, 2-(3-amino)propylamino-pyrimidin-4-yl, 2-(6-amino)hexylamino-pyrimidin-4-yl, 2-(7-amino)heptylamino-pyrimidin-4-yl, 2-(8-amino)octylamino-pyrimidin-4-yl, 2-hydroxylamino-pyrimidin-4-yl, 2-(2-hydroxy-3-hydroxy)propylamino-pyrimidin-4-yl, 2-guanidino-pyrimidin-4-yl, 2-(morpholin-4-yl)-pyrimidin-4-yl, 2-(di(2-hydroxy)ethyl)amino-pyrimidin-4-yl, 2-(1-methyl)piperidin-4-yl-pyrimidin-4-yl, 2-(1-benzyl)piperidin-4-yl-pyrimidin-4-yl, 2-[(1-hydroxymethyl-2-hydroxy)ethylamino]-pyrimidin-4-yl, 2-methyl-pyrimidin-4-yl, 2-aminocarbonyl-pyrimidin-4-yl, 2-cyano-pyrimidin-4-yl, 2-(piperidin-1-yl)-pyrimidin-4-yl, 2-methylamino-pyrimidin-4-yl, 4-pyrimidinyl, 3-pyrazolyl, 2-methylthio-1-methyl-1,3,4-triazol-5-yl, 2-amino-1,3,5-triazin-4-yl, 2-methoxy-pyrimidin-4-yl, 2-(2-carboxy)ethylamino-pyrimidin-4-yl, 2-carboxymethylamino-pyrimidin-4-yl, 2-(2-hydroxy)ethyloxy-pyrimidin-4-yl, 6-hydroxy-2-amino-pyrimidin-4-yl, 2-(2-amino)ethyloxyethyloxyethyl-pyrimidin-4-yl.

20


25

30


35

Most preferred compounds are compounds 9 (see Table 3), 34 (see Table 2), 58 (see Table 2) and 84 (see Table 3).

In general, compounds of formula (I) wherein L is 4-pyrimidinyl substituted with R^a in position 2, wherein R^a represents hydrogen, amino, optionally substituted C₁₋₆alkyl, optionally substituted mono- or di(C₁₋₁₂alkyl)amino, Het-NH- or Het¹, said compounds being represented by formula (I-a), can be prepared by reacting an intermediate of formula (II) with an intermediate of formula (III) or a suitable salt thereof, such as for example formamidine acetate, guanidine hydrochloride, guanidine carbonate (2:1), guanidine sulphate, guanidine hemisulphate, *N*-(3-hydroxypropyl)guanidine hemisulphate, 1-piperazinecarboximidamide sulfate (2:1) and the like, in the presence of a suitable solvent, such as for example *N,N*-dimethylformamide, dimethylsulphoxide, an alcohol, such as for example ethanol, 2-ethoxyethanol, 2-methoxyethanol and the like, a suitable base, e.g. sodium methanolate (sodium methoxide), sodium ethanolate (sodium ethoxide), sodium hydride and the like. Sodium in the presence of a suitable alcohol may also be used. The reaction may be performed at elevated temperature.




Compounds of formula (I-a) wherein Z represents aminocarbonyl, said compounds being represented by formula (I-a-1), can be prepared by reacting an intermediate of formula (II') with an intermediate of formula (III) or a suitable salt thereof, such as for example formamidine acetate, guanidine hydrochloride, guanidine carbonate (2:1), guanidine sulphate, guanidine hemisulphate, *N*-(3-hydroxypropyl)guanidine hemisulphate, 1-piperazinecarboximidamide sulfate (2:1) and the like, in the presence of a suitable solvent, such as for example *N,N*-dimethylformamide, dimethylsulphoxide, an alcohol, such as for example ethanol, 2-ethoxyethanol, 2-methoxyethanol and the like, a suitable base, e.g. sodium methanolate (sodium methoxide), sodium hydride and the like. Sodium in the presence of a suitable alcohol may also be used. The reaction may be performed at elevated temperature.




-27-

Compounds of formula (I) wherein L is 4-pyrimidinyl substituted with R^b in position 2, wherein R^b represents $-NH_2$, Het^1-NH- ; Het^1 ; $-NH-C(=NH)-N(R^2)_2$; C_{1-12} alkyloxy optionally substituted with one, two or three hydroxy groups; optionally substituted mono-or di(C_{1-12} alkyl)amino, in particular unsubstituted mono-or di(C_{1-12} alkyl)amino or mono-or di(C_{1-12} alkyl)amino wherein C_{1-12} alkyl is substituted with one, two or three substituents selected from hydroxy, carboxyl, amino, amino C_{1-4} alkyloxy C_{1-4} alkyloxy, said compounds being represented by formula (I-b), can be prepared by reacting an intermediate of formula (IV) with an intermediate of formula (V) optionally at elevated temperature and optionally in the presence of a suitable solvent, such as for example *N,N*-dimethylformamide, dimethylsulphoxide, tetrahydrofuran, an alcohol, e.g. 2-propanol, methanol, methanol/sodium methoxide and the like, and optionally in the presence of a suitable base, such as for example disodium carbonate, in order to form the corresponding salt wherever possible, or a suitable acid, such as for example hydrochloric acid or acetic acid and the like, and optionally in the presence of sodium hydride, for instance when $H-R^b$ represents hydroxy C_{1-12} alkyloxy, and optionally under pressure.




An analogous reaction can be performed to convert an intermediate of formula (IV') into a compound of formula (I'-b).




Compounds of formula (I) wherein L is 4-pyrimidinyl substituted with cyano in position 2, said compounds being represented by formula (I-c), can be prepared by reacting an intermediate of formula (IV) with a suitable cyanide salt, such as for example potassium cyanide, in the presence of a suitable solvent, such as for example *N,N*-dimethylformamide. The reaction may be performed at elevated temperature.


-28-




- Compounds of formula (I) wherein L is 4-pyrimidinyl substituted with hydroxy in position 2, said compounds being represented by formula (I-d), may be prepared by reacting an intermediate of formula (IV) with a suitable hydroxyde base, such as for example sodium hydroxide, in the presence of a suitable solvent, such as for example water, tetrahydrofuran.



- Compounds of formula (I) wherein L is 4-pyrimidinyl substituted with $\text{CH}_3\text{-S-}$ in position 2, said compounds being represented by formula (I-e), can be prepared by reacting an intermediate of formula (II) with thiourea in the presence of a suitable solvent, such as for example an alcohol, e.g. ethanol and the like, a suitable alcoholate, e.g. sodium ethanolate (sodium ethoxide) and the like, dimethyl sulphate, and a suitable base, such as for example sodium hydroxide.




- Compounds of formula (I) wherein L is 4-pyrimidinyl substituted with R^a in position 2 and $\text{CH}_3\text{-S-}$ in position 6, said compounds being represented by formula (I'-e), can be prepared by reacting an intermediate of formula (XXXV) with an intermediate of formula (III) in the presence of a suitable solvent, such as for example *N,N*-dimethylformamide. The reaction may be performed at elevated temperature.




- Compounds of formula (I) wherein L is 3-pyrazolyl, said compounds being represented by formula (I-f), can be prepared by reacting an intermediate of formula (II) with

-29-


hydrazine monohydrate in the presence of a suitable acid, such as for example acetic acid.



- Compounds of formula (I) wherein L is triazolyl substituted with mercapto, said compounds being represented by formula (I-g), can be prepared by reacting an intermediate of formula (VII) with an intermediate of formula (VIII) wherein R^c represents hydrogen or C₁₋₆alkyl, in the presence of a suitable base, such as for example 1,8-diazabicyclo[5,4,0]undec-7-ene, and a suitable solvent, such as for example an alcohol, e.g. butanol.




- Compounds of formula (I) wherein L is 4-pyrimidinyl substituted with R^a in position 2 and substituted with hydroxy in position 6, said compounds being represented by formula (I-h), can be prepared by reacting an intermediate of formula (VI) with an intermediate of formula (III) or a suitable salt thereof, such as for example formamidine acetate, guanidine hydrochloride, guanidine carbonate (2:1), guanidine sulphate, guanidine hemisulphate, N-(3-hydroxypropyl)guanidine hemisulphate, 1-piperazinecarboximidamide sulfate (2:1) and the like, in the presence of a suitable solvent, such as for example 2-ethoxyethanol/sodium methoxide. The reaction may be performed at elevated temperature.



Compounds of formula (I) wherein L is 1,3,5-triazin-4-yl substituted with R^a in position 2, said compounds being represented by formula (I-i), can be prepared by reacting an intermediate of formula (XXX) with an intermediate of formula (III) or a

suitable salt thereof, such as for example formamidine acetate, guanidine hydrochloride, guanidine carbonate (2:1), guanidine sulphate, guanidine hemisulphate, *N*-(3-hydroxypropyl)guanidine hemisulphate, 1-piperazinecarboximidamide sulfate (2:1) and the like, in the presence of 1,1-dimethoxy-*N,N*-dimethyl-methanamine and in the presence of a suitable solvent, such as for example methanol/sodium methoxide.



Compounds of formula (I) may be converted into each other following art-known functional group transformation reactions, comprising those described hereinafter.

- 10 The compounds of formula (I) may be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth
- 15 alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboxy-peroxy acid or halo substituted benzenecarboxy-peroxy acid, e.g. 3-chlorobenzenecarboxy-peroxy acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g. t.butyl hydro-peroxide. Suitable solvents are, for example, water, lower alcohols, e.g.
- 20 ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

- Compounds of formula (I) wherein L is substituted with amino may be converted into a compound of formula (I) wherein L is substituted with C₁₋₆alkylcarbonylamino by
- 25 reaction with a C₁₋₆alkylcarbonyl chloride in a suitable solvent, such as for example pyridine.


Compounds of formula (I) wherein Q or L is substituted with cyano may be converted into a compound of formula (I), wherein Q or L is substituted with carboxyl by reaction with a suitable acid, such as concentrated hydrochloric acid, in the presence of a suitable reaction-inert solvent, e.g. water.

- 5 Compounds of formula (I) wherein Q or L is substituted with cyano may also be converted into a compound of formula (I), wherein Q or L is substituted with aminocarbonyl by reaction with a suitable acid, such as for example sulphuric acid, in the presence of water.
- 10 Compounds of formula (I), wherein L is substituted with C₁₋₆alkyl-C(=O)-NH-, may be converted into a compound of formula (I), wherein L is substituted with amino, by reaction with a suitable acid, such as for example hydrobromic acid and the like, in the presence of a suitable solvent, such as water.
- 15 Compounds of formula (I) wherein L is substituted with mercapto can be converted into a compound of formula (I) wherein L is substituted with C₁₋₆alkylthio by reaction with a suitable alkylating agent, such as for example C₁₋₆alkyl-I, e.g. CH₃-I, in the presence of a suitable solvent, such as for example an alcohol, e.g. ethanol.
- 20 Compounds of formula (I) wherein L is substituted with fluoro may be prepared from a compound of formula (I) wherein L is not substituted with fluoro by reaction with select fluor in the presence of 2,6-lutidine and a suitable solvent, such as for example *N,N*-dimethylformamide. Compounds of formula (I) wherein L is substituted with chloro or bromo may be prepared from a compound of formula (I) wherein L is not
- 25 substituted with chloro or bromo by reaction with *N*-chlorosuccinimide or *N*-bromosuccinimide in the presence of a suitable solvent, such as for example carbon tetrachloride.


- In the following paragraphs, there are described several methods of preparing the
- 30 intermediates in the foregoing preparations. A number of intermediates and starting materials are commercially available or are known compounds which may be prepared according to conventional reaction procedures generally known in the art.

- Intermediates of formula (II) can be prepared by reacting an intermediate of formula
- 35 (IX) with (CH₃)₂N-CH(OCH₃)₂ at elevated temperature, optionally in the presence of a suitable solvent, such as for example *N,N*-dimethylformamide or toluene.


-32-




Intermediates of formula (IX) wherein Z represents C_{1-6} alkyl or C_{1-6} alkyloxycarbonyl, said Z being represented by Z^a and said intermediates being represented by formula (IX-a), can be prepared by reacting an intermediate of formula (X) with an intermediate of formula (XI) wherein W_1 represents a suitable leaving group, such as halo, for example chloro, bromo, and Z^a represents C_{1-6} alkyl or C_{1-6} alkyloxycarbonyl as described above, in the presence of a suitable solvent, such as for example an alcohol, e.g. methanol, ethanol and the like, at elevated temperature.



Intermediates of formula (IX) can also be prepared by oxidizing an intermediate of formula (XII) in the presence of a suitable oxidation reagent, such as for example pyridinium chlorochromate, in the presence of a suitable solvent, such as for example 1,2-dichloroethane and at elevated temperature.




Intermediates of formula (XII) can be prepared by reacting an intermediate of formula (XIII) with CH_3MgCl in the presence of a suitable solvent, such as for example tetrahydrofuran.




Intermediates of formula (XIII) can be prepared by reacting an intermediate of formula (XIV) wherein W_2 represents a suitable leaving group, such as for example halo, e.g. chloro and the like, with an intermediate of formula (XV) in the presence of a suitable

catalyst, such as for example palladiumtetrakis(triphenylphosphine), and a suitable solvent, such as for example tetrahydrofuran and a suitable salt, such as for example disodium carbonate in water.




- 5 Intermediates of formula (IX) wherein Z represents chloro, said intermediates being represented by formula (IX-b), or intermediates of formula (XIII) wherein Z represents chloro, said intermediates being represented by formula (XIII-a), can be prepared by reacting an intermediate of formula (XVI) with *N,N*-dimethylformamide in the presence of POCl₃ at elevated temperature




10

Intermediates of formula (XVI) can be prepared by reacting an intermediate of formula (X) with Cl-CH₂-C(=O)-Cl at elevated temperature.



15


Intermediates of formula (XIII-a) wherein Q represents 1-pyrrolyl, said intermediates being represented by formula (XIII-a-1), can be prepared by reacting 2-amino-4-chloro-5-thiazolecarboxaldehyde (CAS 76874-79-8) with tetrahydro-2,5-dimethoxy-furan (CAS 696-59-3) in the presence of a suitable acid, such as for example acetic acid.




20

In an analogous manner can 2-amino-4-methyl-5-thiazolyl-ethanone (CAS 106012-40-2) react with tetrahydro-2,5-dimethoxy-furan in the presence of a suitable acid, such as for example acetic acid, to form 2-pyrrol-1-yl-4-methyl-5-thiazolyl-ethanone.


-34-




Intermediates of formula (XIII-a) wherein Q represents 1-pyrazolyl, said intermediates being represented by formula (XIII-a-2), can be prepared by reacting intermediate (XXXI) with an acid, such as for example acetic acid, in the presence of a suitable solvent, such as for example an alcohol, e.g. methanol and the like, at elevated temperature.



Intermediates of formula (XXXI) can be prepared by reacting an intermediate of formula (XXXII) with pyrazole in the presence of a suitable solvent, such as for example *N,N*-dimethylformamide and in the presence of sodium hydride.




Intermediates of formula (XXXII) can be prepared by reacting an intermediate of formula (XXXIII) with ethan-1,2-diol in the presence of a suitable acid, such as 4-toluenesulphonic acid, and a suitable solvent, such as toluene.




Intermediates of formula (II') can be prepared by reacting an intermediate of formula (XVII) with $(\text{CH}_3)_2\text{N-CH}(\text{OCH}_3)_2$ at elevated temperature.

-35-




Intermediates of formula (XVII) can be prepared by reacting an intermediate of formula (XVIII) with NH₃ in the presence of a suitable solvent, such as for example an alcohol, e.g. methanol and the like.




5

Intermediates of formula (XVIII) can be prepared by reacting an intermediate of formula (X) with CH₃-C(=O)-CHCl-C(=O)-C(=O)-O-CH₂CH₃ at elevated temperature in the presence of a suitable solvent, such as for example an alcohol, e.g. ethanol and the like.



10


Intermediates of formula (IV) can be prepared by reacting a compound of formula (I-e) with a suitable oxidation reagent, such as for example 3-chloroperoxybenzoic acid, in the presence of a suitable solvent, such as for example methylene chloride.




15

Intermediates of formula (IV') can be prepared by reacting a compound of formula (I'-e) with a suitable oxidation reagent, such as for example 3-chloroperoxybenzoic acid, in the presence of a suitable solvent, such as for example chloroform, and a suitable base, such as for example disodium carbonate and sodium metabisulphite.


-36-




Intermediates of formula (VI) can be prepared by reacting an intermediate of formula (IX) with $\text{CH}_3\text{-C(=O)-O-C}_{1-4}\text{alkyl}$ at elevated temperature in the presence of a suitable solvent, such as for example tetrahydrofuran, and in the presence of sodium hydride.



Intermediates of formula (IX) wherein Z represents chloro, said intermediates being represented by formula (IX-b), can be converted into an intermediate of formula (IX) wherein Z represents $\text{C}_{1-4}\text{alkyloxy}$, said intermediates being represented by formula (IX-c), by reaction at elevated temperature with a suitable alcoholate, such as for example sodium methanolate (sodium methoxide) in the presence of a suitable solvent, such as for example an alcohol, e.g. methanol.




Intermediates of formula (VII) may be prepared by reacting an intermediate of formula (IXX) with hydrazine hydrate.




Intermediates of formula (XXX) can be prepared by reacting an intermediate of formula (XXXIV) with a suitable anhydride, such as for example trifluoroacetic anhydride.


-37-



Intermediates of formula (XXXIV) can be prepared by reacting an intermediate of formula (XXXV) with ammonia in the presence of oxalyl chloride and a suitable solvent, such as for example methylene chloride and *N,N*-dimethylformamide.




Intermediates of formula (XXXVI) can be prepared by reacting an intermediate of formula (IX) with carbon disulphide and methyl iodide in the presence of a suitable solvent, such as for example tetrahydrofuran, and a suitable base, such as for example potassium *tert*.-butoxide.



The intermediates of formula (IX) wherein Q represents 4-trifluoromethylphenyl and Z represents halo or C_{1-4} alkyl, said Z being represented by formula Z^b , and said intermediates being represented by formula (IX-c), are novel and also form part of the present invention.

Therefore, the present invention also relates to a compound of formula




wherein Z^b represents halo or C_{1-4} alkyl.

Preferred compounds of formula (IX-c) are those compounds wherein Z^b represents halo, in particular chloro.

The intermediates of formula (XIII) wherein Q represents 4-trifluoromethylphenyl and Z represents halo or C_{1-4} alkyl, said Z being represented by formula Z^b , and said

intermediates being represented by formula (XIII-b), are novel and also form part of the present invention.

Therefore, the present invention also relates to a compound of formula



wherein Z^b represents halo or C₁₋₄alkyl.

Preferred compounds of formula (XIII-b) are those compounds wherein Z^b represents halo, in particular chloro.

10 The compounds of the present invention show cytokine production modulating activity, in particular cytokine production inhibitory activity, more in particular proinflammatory cytokine production inhibitory activity. A cytokine is any secreted polypeptide that affects the function of other cells by modulating interactions between

15 cells in the immune or inflammatory response. Examples of cytokines include Interleukin-1 (IL-1) up to Interleukin-23 (IL-23), Tumor Necrosis Factor-alpha (TNF-α), Tumor Necrosis Factor-beta (TNF-β). The present compounds also show inhibitory activity on the production of chemotactic cytokines or chemokines responsible for trafficking and activation of leucocytes. A chemokine production

20 inhibited by the compounds of formula (I) is MCP-1 production (Monocyte Chemotactic Protein 1).

The cytokine production specifically inhibited by the compounds of formula (I) is TNF-α and/or Interleukin-12 (IL-12) production.

25 TNF-α is primarily produced by monocytes, macrophages, T and B lymphocytes, neutrophils, mast cells, tumour cells, fibroblasts, keratinocytes, astrocytes, microglial cells, smooth muscle cells and others. This proinflammatory cytokine is established at the pinnacle of proinflammatory cascades; it exerts a key role in the cytokine network

30 with regard to the pathogenesis of many infectious, inflammatory and autoimmune diseases. Excessive or unregulated TNF-α production is implicated in mediating or exacerbating a number of diseases including rheumatoid arthritis, rheumatoid spondylitis, spondyloarthropathies, systemic lupus erythematosus, osteoarthritis, gouty arthritis, juvenile arthritis and other arthritic conditions, polychondritis, sclerodoma,

Wegener granulomatosis, dermatomyositis, Steven-Johnson syndrome, idiopathic sprue, endocrine ophthalmopathy, Grave's disease, alveolitis, chronic hypersensitivity pneumonitis, primary billiary cirrhosis, uveitis, keratoconjunctivitis sicca and vernal keratoconjunctivitis, allergic rhinitis, pemphigus, eosinophilia, Loffler's syndrome, eosinophilic pneumonia, parasitic infestation, bronchopulmonary aspergillosis, polyarteritis nodosa, eosinophilic granuloma, eosinophil-related disorders affecting the airways occasioned by drug-reaction, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, cerebral malaria, adult respiratory distress syndrome, bronchitis (acute, arachidic, catarrhal, chronic, croupus, phthinoid bronchitis), chronic obstructive airway or pulmonary disease, pulmonary fibrosis, pneumoconiosis (aluminosis, anthracosis, asbestosis, chalicocis, ptilosis, siderosis, silicosis, tobaccosis, byssionosis), tuberculosis, silicosis, exacerbation of airways hyperreactivity to other drug therapy (e.g. aspirin or β -agonist therapy), pulmonary sarcoidosis, bone resorption diseases, meningitis, reperfusion injury, graft versus host reaction, allograft rejections, transplant rejections, fever and myalgias due to infection, such as influenza, cachexia (consequential to, e.g. bacterial, viral or parasitic, infection or to deprivation or deterioration of humoral or other organic function, or secondary to malignancy; malarial and vernal cachexia; cachexia resulting from dysfunction of the pituitary, thyroid or thymus glands as well as uremic cachexia; cachexia secondary to acquired immune deficiency syndrome (AIDS)), AIDS, ARC (AIDS related complex), diabetes, cancer, angiogenesis, lymphoma, Kawasaki syndrome, Behçet's syndrome, aphthous ulceration, skin-related disorders such as psoriasis, eczema, burns, dermatitis, keloid formation, scar tissue formation, erythema nodosum leprosum, Crohn's disease, ulcerative colitis, inflammatory bowel disease, irritable bowel syndrome, pyresis, asthma (intrinsic, extrinsic, allergic, non-atopic, exercise induced and occupational and bacterial infection induced asthma), wheezy infant syndrome, multiple sclerosis, Parkinson's disease, pancreatitis, cardiac disease, congestive heart failure, myocardial infarction, acute liver failure, glomerulonephritis, therapy-associated syndromes comprising Jarisch-Herxheimer reaction, and syndromes associated with IL-2 infusion, anti-CD3 antibody infusion, hemodialysis, yellow fever vaccination.

TNF- α has also been shown to activate HIV (Human Immune deficiency Virus) replication in monocytes and/or macrophages. Therefore, inhibition of TNF- α production or activity aids in limiting HIV progression. TNF- α also plays a role in other viral infections, such as Hepatitis C, CMV (cytomegalovirus), influenza and herpes virus infections, including herpes simplex virus type-1, herpes simplex virus type-2, varicella-zoster virus, Epstein-Barr virus, human herpes virus-6, -7 and -8, pseudorabies and rhinotracheitis.

IL-12 is produced primarily by monocytes, macrophages and dendritic cells in response to bacteria, bacterial products (lipopolysaccharide) and immune signals. The production of IL-12 is regulated by other cytokines and endogenous mediators produced during inflammatory and immunological responses. IL-12 plays a central role in the immune system. Evidence obtained from animal models and human diseases suggests that inappropriate and protracted production of IL-12 and the ability of IL-12 to induce the generation of T helper 1 cell type responses may be instrumental in the development and maintenance of chronic inflammatory diseases, such as rheumatoid arthritis, collagen induced arthritis, allergic encephalitis, colitis, inflammatory bowel disease, Crohn's disease and multiple sclerosis, and in the triggering of autoimmune disorders, such as diabetes, or graft versus host diseases, shock or musculoskeletal and connective tissue diseases. The adverse effects also include anemia (haemolytic, aplastic, pure red cell, idiopathic thrombocytopenia), neutropenia, lymphopenia, hepatosplenomegaly with mononuclear cell infiltration and pulmonary edema with interstitial cell infiltrates. Excessive IL-12 production may accelerate the inflammatory progress of a disease, or the onset of the disease, such as rheumatoid arthritis, or it may also augment the disease severity.

Inhibition of TNF- α and/or IL-12 production by the compounds of formula (I) might offer an interesting, potentially less toxic alternative to non-specific immunosuppression (e.g. corticosteroids) in the treatment of chronic inflammatory and autoimmune diseases. The combined modulation of TNF- α and IL-12 production may ameliorate the treated disease to a greater extent than mono-therapy. The therapeutic effect of combining the suppression of both the immune and the inflammatory arm of a disease may provide additional clinical benefits. The present compounds are also indicated for use as co-therapeutic agents for use in conjunction with immunosuppressive and/or anti-inflammatory drugs, e.g. as potentiators of the therapeutic activity of said drugs, to reduce required dosaging or thus also potential side effects of said drugs. Immunosuppressive and/or anti-inflammatory drugs include for example cyclopeptide, cyclopeptolide or macrolide immunosuppressive or anti-inflammatory drugs, such as drugs belonging to the cyclosporin class, e.g. cyclosporine A or G, tacrolimus substances, ascomycin, rapamycin, glucocorticosteroid drugs, e.g. budesonide, beclamethasone, fluticasone, mometasone.

35

The compounds of formula (I) are useful in preventing or treating cytokine mediated diseases, and as such, inhibit, suppress or antagonize the production or activity of proinflammatory cytokines, such as TNF- α and/or IL-12.

- 5 Disorders mediated through TNF- α and/or IL-12 refers to any and all disorders and disease states in which TNF- α and/or IL-12 play a role, either by the cytokine itself, or by the cytokine causing another cytokine, such as for example IL-1 or IL-6, or a certain mediator to be released.
- 10 Due to their cytokine production inhibitory activity, in particular their proinflammatory cytokine production inhibitory activity, more in particular their TNF- α and/or IL-12 inhibitory activity, even more in particular their IL-12 inhibitory activity, the compounds of formula (I), their N-oxides, pharmaceutically acceptable addition salts, quaternary amines and stereochemically isomeric forms are useful in the treatment or
- 15 prevention of diseases or conditions mediated through cytokines, in particular diseases or conditions related to excessive or unregulated production of proinflammatory cytokines, such as TNF- α and/or IL-12, comprising inflammatory diseases or autoimmune diseases. Diseases or conditions related to an excessive or unregulated production of proinflammatory cytokines comprise rheumatoid arthritis, rheumatoid
- 20 spondylitis, spondyloarthropathies, systemic lupus erythematosus, osteoarthritis, gouty arthritis, juvenile arthritis and other arthritic conditions, polychondritis, sclerodoma, Wegener granulomatosis, dermatomyositis, Steven-Johnson syndrome, idiopathic sprue, endocrine ophthalmopathy, Graves' disease, alveolitis, chronic hypersensitivity pneumonitis, primary billiary cirrhosis, uveitis, keratoconjunctivitis sicca and vernal
- 25 keratoconjunctivitis, allergic rhinitis, pemphigus, eosinophilia, Loffler's syndrome, eosinophilic pneumonia, parasitic infestation, bronchopulmonary aspergillosis, polyarteritis nodosa, eosinophilic granuloma, eosinophil-related disorders affecting the airways occasioned by drug-reaction, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, cerebral malaria, adult respiratory distress
- 30 syndrome, bronchitis (acute, arachidic, catarrhal, chronic, croupus, phthinoid bronchitis), chronic obstructive airway or pulmonary disease, pulmonary fibrosis, tuberculosis, pneumoconiosis (aluminosis, anthracosis, asbestosis, chalicocis, ptilosis, siderosis, silicosis, tobaccosis, byssionosis), exacerbation of airways hyperreactivity to other drug therapy (e.g. aspirin or β -agonist therapy), silicosis, pulmonary sarcoidosis,
- 35 bone resorption diseases, meningitis, allergic encephalitis, reperfusion injury, graft versus host reaction, allograft rejections, transplant rejections, musculoskeletal and connective tissue diseases, fever and myalgias due to infection, such as influenza,

cachexia (consequential to, e.g. bacterial, viral or parasitic, infection or to deprivation or deterioration of humoral or other organic function, or secondary to malignancy; malarial and vermal cachexia; cachexia resulting from dysfunction of the pituitary, thyroid or thymus glands as well as uremic cachexia; cachexia secondary to acquired
5 immune deficiency syndrome (AIDS)), AIDS, ARC (AIDS related complex), diabetes, cancer, angiogenesis, lymphoma, Kawasaki syndrome, Behçet's syndrome, aphthous ulceration, skin-related disorders such as psoriasis, eczema, burns, dermatitis, keloid formation, scar tissue formation, erythema nodosum leprosum, Crohn's disease, ulcerative colitis, inflammatory bowel disease, irritable bowel syndrome, pyresis,
10 asthma (intrinsic, extrinsic, allergic, non-atopic, exercise induced and occupational and bacterial infection induced asthma), wheezy infant syndrome, multiple sclerosis, Parkinson's disease, pancreatitis, cardiac disease, congestive heart failure, myocardial infarction, acute liver failure, glomerulonephritis, therapy-associated syndromes comprising Jarisch-Herxheimer reaction, and syndromes associated with IL-2 infusion,
15 anti-CD3 antibody infusion, hemodialysis, yellow fever vaccination, HIV or other viral infections, such as Hepatitis C, CMV, influenza and herpes virus infections, pseudorabies and rhinotracheitis, angiofollicular lymphoid hyperplasia, anemia (haemolytic, aplastic, pure red cell, idiopathic thrombocytopenia), neutropenia, lymphopenia, hepatosplenomegaly with mononuclear cell infiltration and pulmonary
20 edema with interstitial cell infiltrates; or to prevent these diseases. In particular, the compounds of formula (I) can be used to treat rheumatoid arthritis, Crohn's disease, irritable bowel disease, colitis, psoriasis or multiple sclerosis.

The cytokine production inhibitory activity of the compounds of formula (I) such as the
25 inhibition of TNF- α and/or IL-12 production, may be demonstrated in the *in vitro* test "Inhibition of cytokine production in human whole blood cultures". Suitable *in vivo* tests are "Determination of cytokine in serum of LPS (lipopolysaccharide) and anti-CD3 challenged mice", "Inhibition of LPS-galactosamine induced shock in mice", "Inhibition of collagen induced arthritis in mice".

30 The compounds of formula (I) may also inhibit Interleukin-6 (IL-6).

The present compounds may also act as intermediates for the preparation of further thiazolyl derivatives.

35 In view of the above described pharmacological properties, the compounds of formula (I) or any subgroup thereof, their N-oxides, pharmaceutically acceptable addition salts,

quaternary amines and stereochemically isomeric forms, may be used as a medicine. In particular, the present compounds can be used for the manufacture of a medicament for treating or preventing diseases mediated through cytokines, more in particular diseases mediated through TNF- α and/or IL-12, such as inflammatory and auto-immune diseases.

In view of the utility of the compounds of formula (I), there is provided a method of treating warm-blooded animals, including humans, suffering from or a method of preventing warm-blooded animals, including humans, to suffer from diseases mediated through cytokines, in particular mediated through TNF- α and/or IL-12, such as inflammatory and auto-immune diseases. Said methods comprise the administration, preferably oral administration, of an effective amount of a compound of formula (I), a *N*-oxide form, a pharmaceutically acceptable addition salt, a quaternary amine or a possible stereoisomeric form thereof, to warm-blooded animals, including humans.

The present invention also provides compositions for preventing or treating diseases mediated through cytokines, in particular TNF- α and/or IL-12 comprising a therapeutically effective amount of a compound of formula (I) and a pharmaceutically acceptable carrier or diluent.

The compounds of the present invention or any subgroup thereof may be formulated into various pharmaceutical forms for administration purposes. As appropriate compositions there may be cited all compositions usually employed for systemically administering drugs. To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compound, optionally in addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirable in unitary dosage form suitable, particularly, for administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs, emulsions and solutions; or solid carriers such as starches, sugars, kaolin, diluents, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules, and tablets. Because of their ease in administration, tablets and capsules represent the most

advantageous oral dosage unit forms, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not introduce a significant deleterious effect on the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on, as an ointment. The compounds of the present invention may also be administered via inhalation or insufflation by means of methods and formulations employed in the art for administration via this way. Thus, in general the compounds of the present invention may be administered to the lungs in the form of a solution, a suspension or a dry powder. Any system developed for the delivery of solutions, suspensions or dry powders via oral or nasal inhalation or insufflation are suitable for the administration of the present compounds.

To aid solubility of the compounds of formula (I), suitable ingredients, e.g. cyclodextrins, may be included in the compositions. Appropriate cyclodextrins are α -, β -, γ -cyclodextrins or ethers and mixed ethers thereof wherein one or more of the hydroxy groups of the anhydroglucose units of the cyclodextrin are substituted with C_{1-6} alkyl, particularly methyl, ethyl or isopropyl, e.g. randomly methylated β -CD; hydroxy C_{1-6} alkyl, particularly hydroxyethyl, hydroxy-propyl or hydroxybutyl; carboxy C_{1-6} alkyl, particularly carboxymethyl or carboxy-ethyl; C_{1-6} alkylcarbonyl, particularly acetyl. Especially noteworthy as complexants and/or solubilizers are β -CD, randomly methylated β -CD, 2,6-dimethyl- β -CD, 2-hydroxyethyl- β -CD, 2-hydroxyethyl- γ -CD, 2-hydroxypropyl- γ -CD and (2-carboxymethoxy)propyl- β -CD, and in particular 2-hydroxypropyl- β -CD (2-HP- β -CD).

The term mixed ether denotes cyclodextrin derivatives wherein at least two cyclodextrin hydroxy groups are etherified with different groups such as, for example, hydroxy-propyl and hydroxyethyl.

- 5 The average molar substitution (M.S.) is used as a measure of the average number of moles of alkoxy units per mole of anhydroglucose. The average substitution degree (D.S.) refers to the average number of substituted hydroxyls per anhydroglucose unit. The M.S. and D.S. value can be determined by various analytical techniques such as nuclear magnetic resonance (NMR), mass spectrometry (MS) and infrared
10 spectroscopy (IR). Depending on the technique used, slightly different values may be obtained for one given cyclodextrin derivative. Preferably, as measured by mass spectrometry, the M.S. ranges from 0.125 to 10 and the D.S. ranges from 0.125 to 3.

- Other suitable compositions for oral or rectal administration comprise particles
15 consisting of a solid dispersion comprising a compound of formula (I) and one or more appropriate pharmaceutically acceptable water-soluble polymers.

- The term "a solid dispersion" used hereinafter defines a system in a solid state (as opposed to a liquid or gaseous state) comprising at least two components, in casu the
20 compound of formula (I) and the water-soluble polymer, wherein one component is dispersed more or less evenly throughout the other component or components (in case additional pharmaceutically acceptable formulating agents, generally known in the art, are included, such as plasticizers, preservatives and the like). When said dispersion of the components is such that the system is chemically and physically uniform or
25 homogenous throughout or consists of one phase as defined in thermo-dynamics, such a solid dispersion will be called "a solid solution". Solid solutions are preferred physical systems because the components therein are usually readily bioavailable to the organisms to which they are administered. This advantage can probably be explained by the ease with which said solid solutions can form liquid solutions when contacted
30 with a liquid medium such as the gastro-intestinal juices. The ease of dissolution may be attributed at least in part to the fact that the energy required for dissolution of the components from a solid solution is less than that required for the dissolution of components from a crystalline or microcrystalline solid phase.

- 35 The term "a solid dispersion" also comprises dispersions which are less homogenous throughout than solid solutions. Such dispersions are not chemically and physically uniform throughout or comprise more than one phase. For example, the term "a solid

dispersion" also relates to a system having domains or small regions wherein amorphous, microcrystalline or crystalline compound of formula (I), or amorphous, microcrystalline or crystalline water-soluble polymer, or both, are dispersed more or less evenly in another phase comprising water-soluble polymer, or compound of
5 formula (I), or a solid solution comprising compound of formula (I) and water-soluble polymer. Said domains are regions within the solid dispersion distinctively marked by some physical feature, small in size, and evenly and randomly distributed throughout the solid dispersion.

10 Various techniques exist for preparing solid dispersions including melt-extrusion, spray-drying and solution-evaporation.

The solution-evaporation process comprises the following steps :

a) dissolving the compound of formula (I) and the water-soluble polymer in an
15 appropriate solvent, optionally at elevated temperatures;
b) heating the solution resulting under point a), optionally under vacuum, until the solvent is evaporated. The solution may also be poured onto a large surface so as to form a thin film, and evaporating the solvent therefrom.

20 In the spray-drying technique, the two components are also dissolved in an appropriate solvent and the resulting solution is then sprayed through the nozzle of a spray dryer followed by evaporating the solvent from the resulting droplets at elevated temperatures.

25 The preferred technique for preparing solid dispersions is the melt-extrusion process comprising the following steps :

a) mixing a compound of formula (I) and an appropriate water-soluble polymer,
b) optionally blending additives with the thus obtained mixture,
c) heating and compounding the thus obtained blend until one obtains a
30 homogenous melt,
d) forcing the thus obtained melt through one or more nozzles; and
e) cooling the melt till it solidifies.

The terms "melt" and "melting" should be interpreted broadly. These terms not only
35 mean the alteration from a solid state to a liquid state, but can also refer to a transition to a glassy state or a rubbery state, and in which it is possible for one component of the mixture to get embedded more or less homogeneously into the other. In particular

cases, one component will melt and the other component(s) will dissolve in the melt thus forming a solution, which upon cooling may form a solid solution having advantageous dissolution properties.

- 5 After preparing the solid dispersions as described hereinabove, the obtained products can be optionally milled and sieved.

The solid dispersion product may be milled or ground to particles having a particle size of less than 600 μm , preferably less than 400 μm and most preferably less than 125 μm .

10

The particles prepared as described hereinabove can then be formulated by conventional techniques into pharmaceutical dosage forms such as tablets and capsules.

- It will be appreciated that a person of skill in the art will be able to optimize the parameters of the solid dispersion preparation techniques described above, such as the most appropriate solvent, the working temperature, the kind of apparatus being used, the rate of spray-drying, the throughput rate in the melt-extruder

- The water-soluble polymers in the particles are polymers that have an apparent viscosity, when dissolved at 20°C in an aqueous solution at 2 % (w/v), of 1 to 5000 mPa.s more preferably of 1 to 700 mPa.s, and most preferred of 1 to 100 mPa.s. For example, suitable water-soluble polymers include alkylcelluloses, hydroxyalkylcelluloses, hydroxyalkyl alkylcelluloses, carboxyalkylcelluloses, alkali metal salts of carboxyalkylcelluloses, carboxyalkylalkylcelluloses, carboxyalkylcellulose esters, starches, pectines, chitin derivates, di-, oligo- and polysaccharides such as trehalose, alginic acid or alkali metal and ammonium salts thereof, carrageenans, galactomannans, tragacanth, agar-agar, gummi arabicum, guar gummi and xanthan gummi, polyacrylic acids and the salts thereof, polymethacrylic acids and the salts thereof, methacrylate copolymers, polyvinylalcohol, polyvinylpyrrolidone, copolymers of polyvinylpyrrolidone with vinyl acetate, combinations of polyvinylalcohol and polyvinylpyrrolidone, polyalkylene oxides and copolymers of ethylene oxide and propylene oxide. Preferred water-soluble polymers are hydroxypropyl methylcelluloses.

- Also one or more cyclodextrins can be used as water soluble polymer in the preparation of the above-mentioned particles as is disclosed in WO 97/18839. Said cyclodextrins include the pharmaceutically acceptable unsubstituted and substituted cyclodextrins

known in the art, more particularly α , β or γ cyclodextrins or the pharmaceutically acceptable derivatives thereof.

5 Substituted cyclodextrins which can be used to prepare the above described particles include polyethers described in U.S. Patent 3,459,731. Further substituted cyclodextrins are ethers wherein the hydrogen of one or more cyclodextrin hydroxy groups is replaced by C₁-6alkyl, hydroxyC₁-6alkyl, carboxy-C₁-6alkyl or C₁-6alkyloxycarbonylC₁-6alkyl or mixed ethers thereof. In particular such substituted
10 cyclodextrins are ethers wherein the hydrogen of one or more cyclodextrin hydroxy groups is replaced by C₁-3alkyl, hydroxyC₂-4alkyl or carboxyC₁-2alkyl or more in particular by methyl, ethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, carboxy-methyl or carboxyethyl.

Of particular utility are the β -cyclodextrin ethers, e.g. dimethyl- β -cyclodextrin as
15 described in Drugs of the Future, Vol. 9, No. 8, p. 577-578 by M. Nogradi (1984) and polyethers, e.g. hydroxypropyl β -cyclodextrin and hydroxyethyl β -cyclodextrin, being examples. Such an alkyl ether may be a methyl ether with a degree of substitution of about 0.125 to 3, e.g. about 0.3 to 2. Such a hydroxypropyl cyclodextrin may for example be formed from the reaction between β -cyclodextrin and propylene oxide and
20 may have a MS value of about 0.125 to 10, e.g. about 0.3 to 3.

Another type of substituted cyclodextrins is sulfobutylcyclodextrines.

The ratio of the compound of formula (I) over the water soluble polymer may vary
25 widely. For example ratios of 1/100 to 100/1 may be applied. Interesting ratios of the compound of formula (I) over cyclodextrin range from about 1/10 to 10/1. More interesting ratios range from about 1/5 to 5/1.

It may further be convenient to formulate the compounds of formula (I) in the form of
30 nanoparticles which have a surface modifier adsorbed on the surface thereof in an amount sufficient to maintain an effective average particle size of less than 1000 nm. Useful surface modifiers are believed to include those which physically adhere to the surface of the compound of formula (I) but do not chemically bond to said compound.

35 Suitable surface modifiers can preferably be selected from known organic and inorganic pharmaceutical excipients. Such excipients include various polymers, low molecular

weight oligomers, natural products and surfactants. Preferred surface modifiers include nonionic and anionic surfactants.

5 Yet another interesting way of formulating the compounds of formula (I) involves a pharmaceutical composition whereby the compounds of formula (I) are incorporated in hydrophilic polymers and applying this mixture as a coat film over many small beads, thus yielding a composition which can conveniently be manufactured and which is suitable for preparing pharmaceutical dosage forms for oral administration.

10 Said beads comprise a central, rounded or spherical core, a coating film of a hydrophilic polymer and a compound of formula (I) and optionally a seal-coating layer.

Materials suitable for use as cores in the beads are manifold, provided that said materials are pharmaceutically acceptable and have appropriate dimensions and
15 firmness. Examples of such materials are polymers, inorganic substances, organic substances, and saccharides and derivatives thereof.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in unit dosage form for ease of administration and uniformity of dosage.
20 Unit dosage form as used herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such unit dosage forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, suppositories,
25 injectable solutions or suspensions and the like, and segregated multiples thereof.

The present compounds are orally active compounds, and are preferably orally administered.

The exact dosage and frequency of administration depends on the particular compound
30 of formula (I) used, the particular condition being treated, the severity of the condition being treated, the age, weight, sex, extent of disorder and general physical condition of the particular patient as well as other medication the individual may be taking, as is well known to those skilled in the art. Furthermore, it is evident that said effective daily amount may be lowered or increased depending on the response of the treated
35 subject and/or depending on the evaluation of the physician prescribing the compounds of the instant invention.

The compounds of formula (I) may also be used in combination with other conventional anti-inflammatory or immunosuppressive agents, such as steroids, cyclooxygenase-2 inhibitors, non-steroidal-anti-inflammatory drugs, TNF- α antibodies, such as for example acetyl salicylic acid, bufexamac, diclofenac potassium, sulindac, diclofenac sodium, ketorolac trometamol, tolmetine, ibuprofen, naproxen, naproxen sodium, tiaprofen acid, flurbiprofen, mefenamic acid, niflumonic acid, meclofenamate, indomethacin, proglumetacine, ketoprofen, nabumetone, paracetamol, piroxicam, tenoxicam, nimesulide, fenylbutazon, tramadol, beclomethasone dipropionate, betamethasone, beclomethasone, budesonide, fluticasone, mometasone, dexamethasone, hydrocortisone, methylprednisolone, prednisolone, prednisone, triamcinolone, celecoxib, rofecoxib, infliximab, leflunomide, etanercept, CPH 82, methotrexate, sulfasalazine, antilymphocytory immunoglobulines, antithymocytory immunoglobulines, azathioprine, cyclosporine, tacrolimus substances, ascomycin, rapamycin, muromonab-CD3.


Thus, the present invention also relates to the combination of a compound of formula (I) and another anti-inflammatory or immunosuppressive agent. Said combination may be used as a medicine. The present invention also relates to a product containing (a) a compound of formula (I), and (b) another anti-inflammatory or immunosuppressive compound, as a combined preparation for simultaneous, separate or sequential use in the treatment of diseases related to an excessive or unregulated cytokine production. The different drugs may be combined in a single preparation together with pharmaceutically acceptable carriers.

Experimental part

A. Preparation of the intermediate compounds


Example A1A

a) Preparation of intermediate 1a



Palladium (II) acetate (0.0003 mol) and 1,3-bis(diphenylphosphino)-propane (0.0006 mol) were added to a solution of 2-chloro-5-fluoropyridine (0.01 mol) in tetrahydrofuran (100 ml) in an autoclave. Liquid ammonia (0.6 mol) was added, and carbon monoxide at a pressure of 40 atm admitted. The mixture was heated at 150°C for 16 hours. After cooling to room temperature, methanol was added (200 ml), and the mixture stirred for 1 hour. The mixture was filtered, and the residue washed with methanol. The combined filtrates were evaporated to dryness under reduced pressure, the residue triturated under di-isopropylether, and dried under reduced pressure. Yield: 0.56 g of intermediate 1a (40%)


b) Preparation of intermediate 1b



A mixture of 5-chloro-2-pyridinecarboxamide (0.004 mol), phosphorous pentasulphide (0.004 mol), and tetrahydrofuran (25 ml) were heated under reflux for 2 hours. The mixture was cooled to room temperature, filtered, and the residue washed with tetrahydrofuran. The residue was suspended in water (20 ml), and the mixture boiled for 15 minutes. After cooling it was extracted with dichloromethane:methanol 9:1. The phases were separated, the organic layer dried (MgSO₄), and the solvent removed under reduced pressure. Yield: 0.57 g of intermediate 1b (82%).


Example A1B

a) Preparation of intermediate 1



A mixture of 3-chloro-2,4-pentanedione (0.098 mol) and 4-trifluoromethylphenylcarbothioamide (0.098 mol) in ethanol (160 ml) was stirred and refluxed for 16 hours. The mixture was filtered, and the residue washed with ethanol and dried under reduced pressure. Yield: 17 g intermediate 1 (60%) (mp. 87-88°C).


b) Preparation of intermediate 2



A suspension of intermediate 1 (0.035 mol) in 1,1-dimethoxy-*N,N*-dimethylmethanamine (150 ml) was heated at 80°C for 6 hours. The solvent was removed under reduced pressure, and the residue triturated under di-isopropyl ether. The mixture was filtered, the residue washed with di-isopropyl ether and then dried under reduced pressure. Yield: 11.0 g of intermediate 2 (92%) (mp. 147°C).

Example A1C

Preparation of intermediate 2a




Dimethyl carbonate (0.07 mol) was added to suspension of sodium hydride (0.07 mol) in tetrahydrofuran (70 ml), and the mixture heated to 60°C. A solution of intermediate

1 (0.035 mol) in tetrahydrofuran (50 ml) was added dropwise. The reaction mixture was heated under reflux for 45 minutes, cooled to room temperature, and methanol added to destroy any remaining sodium hydride. The mixture was neutralised with acetic acid, and the solvent removed under reduced pressure. The residue was
5 partitioned between ethyl acetate and water, the phases separated, and the aqueous layer extracted twice with ethyl acetate. The combined organic layers were washed with brine, dried (MgSO_4), and the solvent removed under reduced pressure. The resultant oil was triturated under diethyl ether/hexane, the mixture filtered, and the residue washed with hexane. Yield : 10.4 g intermediate 2a (87%).


10 Example A2

a) Preparation of intermediate 3




$\text{Pd(PPh}_3)_4$ (0.002 mol) was dissolved in tetrahydrofuran (120 ml). 2,4-dichloro-5-thiazolecarboxaldehyde (0.02 mol) was added. The reaction mixture was stirred at room temperature for 30 minutes. [4-(trifluoromethyl)phenyl]boronic acid (0.021 mol) and $\text{Na}_2\text{CO}_3/\text{H}_2\text{O}$ (11 g/80 ml) were added. The reaction mixture was
15 stirred and refluxed for 3 hours. The reaction mixture was cooled, water was added. The mixture was extracted with CH_2Cl_2 . The organic layer was separated, dried (MgSO_4), and the solvent evaporated. The residue was triturated under ethanol, washed with ethanol and dried under reduced pressure. Yield: 3,2 g of intermediate 3 (55%).

b) Preparation of intermediate 4



20 Intermediate 3 (0.011 mol) was dissolved in tetrahydrofuran (50 ml) and cooled to -10°C . A 3 M solution of methyl magnesiumchloride in tetrahydrofuran (3.7 ml, 0.011 mol) was added dropwise. The reaction mixture was stirred at room temperature for 3 hours. Water (3 ml) was added, followed by CH_3COOH (1 ml) (exothermic). The reaction mixture was extracted with ethyl acetate, the organic layer was separated,
25 dried, filtered, and the solvent was evaporated. Yield: 3.4 g of intermediate 4 (100%).


c) Preparation of intermediate 5



-53-

Intermediate 4 (0.01 mol) was dissolved in 1,2-dichloroethane (40 ml). Pyridinium chlorochromate (0.02 mol) was added to the reaction mixture. The reaction mixture was stirred at 70°C for 3 hours, cooled to room temperature, poured onto a silica gel plug and eluted with dichloromethane. The eluent was evaporated under reduced pressure, and the residue dried under reduced pressure. Yield: 2.2 g of intermediate 5 (72%) (mp. 178°C).


d) Preparation of intermediate 6



Intermediate 5 (0.004 mol) was dissolved in 1,1-dimethoxy-*N,N*-dimethylmethanamine (50 ml) and stirred at 80°C for 16 hours. The reaction mixture was cooled, water was added, and the mixture filtered. The residue was washed with water and dried under reduced pressure. Yield: 1.1 g of intermediate 6 (76%) (mp. 198°C).


Example A3A

a) Preparation of intermediate 7




2-Amino-4-chloro-5-thiazolecarboxaldehyde (0.012 mol) was dissolved in CH₃COOH (80 ml) and heated. Tetrahydro-2,5-dimethoxyfuran (0.015 mol) was added dropwise to the hot solution. The reaction mixture heated under reflux for 2 hours. The solvent was evaporated. The residue was dried under reduced pressure. Yield : 2.0 g of intermediate 7 (78%).

Intermediate 7 was transformed into intermediate 7'



in a manner analogous to the transformation of intermediate 3 into intermediate 6 as described in Example A2. Yield : 57 % of intermediate 7' (mp. 201°C).

b) Preparation of intermediate 7a




2-Amino-4-methyl-5-acetylthiazole (0.030 mol) was dissolved in CH₃COOH (160 ml) and heated. Tetrahydro-2,5-dimethoxyfuran (0.035 mol) was added dropwise to the hot solution. The reaction mixture was stirred under reflux for 2 hours. The solvent was evaporated. The residue was dried under reduced pressure. Yield : 5.3 g of intermediate 7a (86%).

-54-


Example A3B

a) Preparation of intermediate 7b




2,4-dichloro-5-thiazolecarboxaldehyde (0.027 mol), ethan-1,2-diol, and 4-toluenesulphonic acid were dissolved in toluene (60 ml) and heated under a Dean-Stark trap for 16 hours. The solvent was removed under reduced pressure and the residue chromatographed on silica gel using dichloromethane as eluent. Yield : 2.0 g of intermediate 7b (33%).

b) Preparation of intermediate 7c




Sodium hydride (0.009 mol) was added portionwise to a stirred suspension of pyrazole (0.009 mol) in *N,N*-dimethylformamide (20 ml). Stirring was continued for 1 hour, and then intermediate 7b (0.009 mol) was added. The mixture was stirred for 72 hours, and ice-water added carefully. The mixture was filtered, the residue washed with water and then dried under reduced pressure. Yield : 1.7 g of intermediate 7c (74%).

c) Preparation of intermediate 7d



Intermediate 7c (0.0058 mol) was added to a solution of acetic acid (5 ml) in water (30 ml) and sufficient methanol added to cause dissolution. The solution was heated under reflux for 1 hour, cooled and then filtered. The residue was washed with water, and dried under reduced pressure. Yield : 1.2 g of intermediate 7d (97%).


d) Preparation of intermediate 7e



Intermediate 7d (0.0052 mol) was transformed into intermediate 7e (0.0028 mol) in a manner analogous to the transformation of intermediate 3 into intermediate 6 as described in Example A2. Yield : 0.8 g of intermediate 7e (54%) (mp 232°C).


Example A4

a) Preparation of intermediate 8



4-fluorophenylcarbothioamide (0.05 mol) and chloroacetyl chloride (0.2 mol) were mixed and stirred at room temperature for 10 minutes. The temperature of the reaction mixture was slowly increased to 60°C. After 30 minutes the reaction mixture was cooled to room temperature, and the volatile components were removed under reduced pressure. The residue was dissolved in diethyl ether, hexane was added, and the mixture was filtered. The residue was washed with diethyl ether/hexane (1/4), and then dried under reduced pressure. Yield : 8.9 g of intermediate 8 (77%).


b) Preparation of intermediate 9



- Phosphorous oxychloride (0.44 mol) was added dropwise to *N,N*-dimethylformamide (0.22 mol) at between 0°C and 5°C with rapid stirring. The reaction mixture was stirred for 30 minutes at 0°C and then intermediate 8 (0.044 mol) was added while allowing the reaction mixture to warm up to room temperature. The reaction mixture was heated to 60°C for 1 hour, and then heated under reflux for 90 minutes. Water was cautiously added, the mixture was filtered. The residue was triturated under diethyl ether, and dried under reduced pressure. Yield: 6.8 g of intermediate 9 (64%).

Example A5

a) Preparation of intermediate 11




- 4-trifluoromethylphenylcarbothioamide (0.10 mol) was dissolved in ethanol (150 ml) and ethyl-3-chloroacetopyruvate (0.11 mol) dissolved in ethanol (50 ml) was added. The reaction mixture was stirred under reflux for 90 minutes. The solvent was evaporated under reduced pressure and the residue was partitioned between CH₂Cl₂ and saturated aqueous sodium bicarbonate. The phases were separated and the aqueous phase extracted twice with CH₂Cl₂. The combined organic layers were washed with water and then brine, dried (MgSO₄), and concentrated to dryness. The crude oily

-56-


residue was triturated under hexane, chromatographed on silica gel using ethyl acetate:hexane (1:4) as eluent to give intermediate 11. Yield : 8.2 g of intermediate 11 (24%) (mp. 72-74°C).

b) Preparation of intermediate 12



5 A solution of ammonia in methanol (7 M, 80 ml) was added dropwise to a cooled, stirred suspension of intermediate 11 (0.01 mol) in methanol (30 ml) at 0°C. The reaction mixture was stirred at 0°C for a further 30 minutes and was then stirred at room temperature for 16 hours. The solvent was removed under reduced pressure, methanol added, and the bulk of the solvent removed under reduced pressure. The solution was cooled to 0°C, filtered, and the residue washed with cold methanol and
10 then hexane. The crude product was recrystallised from ethanol:water (3:1). Yield : 1.7 g (54%) of intermediate 12 (mp. 196-200°C).


c) Preparation of intermediate 13



Intermediate 12 (0.002 mol) was mixed with 1,1-dimethoxy-*N,N*-dimethylmethanamine (0.010 mol), and the mixture was heated at 110°C for 40 minutes. After cooling to room temperature the mixture was filtered and the residue was triturated
15 under hexane and separated. Yield: 0.84 g intermediate 13 (95%).

Example A6

Preparation of intermediate 14




Compound 37 (prepared according to B6) (0.035 mol) and 3-chloroperoxybenzoic acid (0.077 mol) were stirred in dichloromethane for 16 hours. The reaction mixture was
20 diluted with dichloromethane (150 ml), and washed with 5% aqueous sodium metabisulphite (3 x 100 ml), and then 5% aqueous sodium carbonate (2 x 100 ml). The reaction mixture was then washed repeatedly with water until the aqueous washings had reached pH 7. The undissolved solids were filtered off. The filtrate was dried (MgSO₄), and evaporated to dryness under reduced pressure to give a residue which

-57-

was combined with the undissolved solids. Yield: 14 g of intermediate 14 (100%) (mp 223°C).



Intermediate 14a



was prepared from compound 91 using the above method for intermediate 14. Yield :
5 74 % of intermediate 14a.

Example A7


Preparation of intermediate 15



(prepared from intermediate 7 analogously to Example A2 b) and c))
(0.0044 mol) was dissolved in methanol (50 ml) and heated until total dissolution. A
10 30wt% solution of sodium methoxide in methanol (0.02 mol) was added. The reaction
mixture was stirred and refluxed for 4 hours and stirred at room temperature overnight.
The solvent was removed under reduced pressure. The residue was washed twice with
water, and dried under reduced pressure. Yield: 0.7 g of intermediate 15 (71%).

15 Example A8


a) Preparation of intermediate 16



Oxalyl chloride (0.039 mol) was added dropwise to a stirred suspension of
4-methyl-2-(4-trifluoromethylphenyl)-5-thiazolecarboxylic acid (0.039 mol) and
N,N-dimethylformamide (1 drop) in dichloromethane (200 ml) at 0°C. The mixture was
allowed to warm to room temperature with stirring over 16 hours. The solvent was
20 removed under reduced pressure, and dichloromethane (100 ml) was added. A solution
of ammonia in methanol (7 M, 30 ml) was added, and the mixture stirred for 8 hours.
The mixture was filtered, the residue washed with dichloromethane and then dried
under reduced pressure. Yield : 10 g of intermediate 16 (90%) (mp 213-216°C).

-58-


b) Preparation of intermediate 17



Trifluoroacetic anhydride (0.0063 mol) was added dropwise to a stirred suspension of intermediate 16 (0.0016 mol) and pyridine (0.010 mol) in 1,4-dioxane at 0°C. Stirring was continued for 30 minutes, the reaction mixture was allowed to warm to room temperature, and the mixture was stirred for a further 16 hours. The solvent was removed under reduced pressure, the residue triturated under water and dried under reduced pressure. Yield : 0.39 g of intermediate 17 (91%) (mp 90°C).

Example A9


Preparation of intermediate 18



A solution of intermediate 1 (0.020 mol) in tetrahydrofuran (25 ml) was added dropwise to a stirred suspension of potassium tert-butoxide (0.040 mol) in tetrahydrofuran (100 ml) at room temperature. A deep red solution formed which was stirred for a further 30 minutes. Carbon disulphide (1.5 ml) was added dropwise, and after 30 minutes methyl iodide (4 ml) was added. The reaction mixture was stirred for 2 hours, then poured into water (1 l), and filtered. The residue was washed with water and dried under reduced pressure. Yield : 4.6 g of intermediate 18 (59%), mp. 164-166 (decomposition).

Example A10

Preparation of intermediate 19




3-chloroperoxybenzoic acid (0.0045 mol) was added portionwise to a stirred suspension of compound 99 (0.0015 mol) in chloroform (10 ml) at room temperature. Stirring was continued for 16 hours. Chloroform (50 ml), and then saturated aqueous sodium metabisulphite (10 ml) were added, and rapid stirring continued for 15 minutes. Saturated aqueous disodium carbonate was added dropwise under gas evolution ceased. The mixture was washed with water and the phases were separated. The organic layer was dried (Na₂SO₄), and the solvent removed under reduced pressure. Yield : 0.60 g of intermediate 19 (97%) (mp.155-158°C).

B. Preparation of the final compounds

Example B1


Preparation of compound 9



Sodium methoxide (0.045 mol) was added to a mixture of diguanidine carbonate
5 (0.023 mol) in 2-ethoxyethanol (300 ml) and the mixture was stirred until a
homogeneous solution was obtained. Intermediate 2 (0.023 mol) was added and the
reaction mixture was heated under reflux for 3 hours. After cooling, water was added
and the mixture was filtered. The residue was washed with water and dried under
reduced pressure. Yield : 5.9 g of compound 9 (76%).

Alternative solvents are ethanol, *N,N*-dimethylformamide or dimethylsulphoxide.


Example B2



A mixture of (0.016 mol) (prepared according to the
15 procedures described in Ex. A1B) and hydrazine monohydrate (0.018 mol) in acetic
acid (20 ml) were stirred under reflux overnight. Boiling water (100 ml) was added to
the hot reaction mixture, and the resultant solution was allowed to cool. The mixture
was filtered and the residue recrystallised from ethanol. Yield : 2.2g of compound 28
(57%).

Example B3

a)




(0.016 mol) (prepared according to the procedures
described in Ex. A1B) was added to a mixture of methylguanidine hydrochloride
25 (0.024 mol) and sodium methoxide (0.026 mol) in *N,N*-dimethylformamide (30 ml).
The mixture was heated at 100 °C for 26 hours. The mixture was diluted with water
(80 ml) and acidified with acetic acid (2 ml). The mixture was filtered. The residue

-60-

was dried and recrystallized from isooctane:toluene 3:1. Yield : 4.4 g of compound 17 (87%).


b)



(0.0073 mol) (prepared according to the procedures

described in Ex. A1B) was added to a mixture of formamidine acetate (0.022 mol) and sodium ethoxide (0.024 mol) in ethanol (20 ml), and the mixture was heated under reflux for 24 hours. A mixture of formamidine acetate (0.012 mol) and sodium ethoxide (0.013 mol) in ethanol (10 ml) was added, and the mixture heated under reflux for a further 24 hours. The mixture was diluted with water (30 ml) and acidified with acetic acid (3 ml). The solvent was removed under reduced pressure. The residue was dried under reduced pressure, and recrystallized from butan-1-ol. Yield : 1.2 g of compound 22 (57%).


c) Preparation of compound 59



Sodium methoxide (0.0014 mol) was added to a mixture of 3-hydroxypropylguanidine hemisulphate (0.0014 mmol) in 2-methoxyethanol. Stirring was continued for 30 minutes, and then intermediate 7' (0.0007 mol) was added. The mixture was stirred at 100 °C for 16 hours, cooled, and the solvent was removed under reduced pressure. The residue was chromatographed on silica gel using dichloromethane as eluent. The residue was triturated under di-isopropylether, filtered, washed with di-isopropylether, and dried under reduced pressure. Yield : 0.013 g of compound 59 (6%).

Example B4

Preparation of compound 35




Sodium methoxide (0.0041 mol) was suspended in 2-ethoxyethanol (6 ml), and acetamidine hydrochloride (0.0041 mol) was added. The mixture was stirred for 30 minutes at room temperature. Intermediate 2 (prepared according to Ex. A1Bb) (0.0018 mol) was added, and the mixture was stirred under reflux for 8 hours. A slurry of sodium methoxide (0.020 mol) and acetamidine hydrochloride (0.020 mol) in

2-ethoxyethanol (2 ml) was added. After stirring under reflux for 1 hour, the reaction mixture was cooled and poured into ice-cold water (70 ml). The mixture was filtered and the residue washed with water. Yield: 0.59 g of compound 35 (98%).

5 Example B5

Preparation of compound 36




Guanidine hydrochloride (0.0054 mol) was added to a solution of sodium methoxide (0.0054 mol) in ethanol (10 ml), and the mixture was stirred for 30 minutes. A solution of intermediate 13 (prepared according to A5c) (0.0018 mol) in ethanol (10 ml) was added, and the reaction mixture refluxed for 1 hour. After cooling, water was added, the mixture was filtered, and the residue washed with ethanol-water. The residue was dried and recrystallized from 2-ethoxyethanol. Yield : 0.23 g of compound 36 (35%) (mp. 286-287°C).

Example B6

Intermediate 2 (0.009 mol) and thiourea (0.010 mol) were added to a solution of potassium hydroxide (0.009 mol) in ethanol (25 ml), and the resulting mixture was refluxed for 5 hours. The mixture was cooled on an ice-bath, filtered, and the residue was washed with diethyl ether. The residue was dried under reduced pressure, and then dissolved in a solution of sodium hydroxide (0.027 mol) in water (40 ml). Dimethyl sulphate (0.018 mol) was added dropwise at room temperature. After 2 hours, water (10 ml) was added and the reaction mixture was extracted with diethyl ether (2 x 40 ml). The organic layers were combined, washed with water (10 ml) and then brine (10 ml), and then dried (MgSO₄). The solvent was removed under reduced pressure. Yield : 2.28 g of compound 37 (69%).

25


Compound 91



was prepared from intermediate 6 using the above method for compound 37. Yield : 52% of compound 91.

Example B7

a) Preparation of compound 38




Intermediate 14 (prepared according to A6) (0.001 mol) was suspended in propane-1,3-diamine (2 ml), and the mixture was heated at 130°C for 15 minutes.

- 5 After cooling, water was added with stirring, and the mixture stood for 16 hours. The mixture was filtered and the residue washed with water. The residue was dried under reduced pressure. Yield : 0.35 g of compound 38 (89%).


- 10 In analogous reactions, a suitable solvent, such as *N,N*-dimethylformamide, may optionally be added if the amine used is not liquid at 20°C.

b) Preparation of compound 84



- 15 Intermediate 14 (prepared according to A6) (0.0005 mol), 3-aminopropionic acid (0.001 mol), and disodium carbonate (0.001 mol) were suspended in dimethylsulphoxide and the mixture was heated to 120°C for 3 hours. The mixture was cooled to room temperature and water (6 ml) was added with stirring. Stirring was continued until crystallisation was complete, the mixture was filtered, and the residue dried under reduced pressure. Yield : 0.2 g of compound 84 (93%).

c) Preparation of compound 83




- 20 A solution of sodium methoxide in methanol (30 wt%, 0.2 ml) was added to a solution of intermediate 14 (prepared according to A6) (0.0002 mol), and methanol (0.8 ml) in tetrahydrofuran (4 ml), and the solution was stirred for 16 hours at room temperature. The solvent was removed under reduced pressure, and the residue stirred in acetonitrile

-63-


(2 ml) and water (4 ml). The mixture was filtered and the residue dried under reduced pressure. Yield : 0.051 g of compound 83 (73%).

d) Preparation of compound 82




5 A solution of sodium hydroxide (1 M, 0.4 ml) was added to a solution of intermediate 14 (prepared according to A6) (0.0002 mol) and water (0.6 ml) in tetrahydrofuran (4 ml). The solution was briefly warmed, and then stirred at room temperature for 72 hours. The solvent was removed under reduced pressure, and the residue stirred in acetonitrile (2 ml), water (2 ml), and aqueous hydrochloric acid (1 M, 0.4 ml). The mixture was filtered and the residue dried under reduced pressure. Yield : 0.058 g of compound 82 (86%).

e) Preparation of compound 40



10 Intermediate 14 (prepared according to A6) (0.0013 mol) was suspended in *N,N*-dimethylformamide (5 ml) and potassium cyanide (0.003 mol) was added. The reaction mixture was heated at 100°C for 15 minutes, cooled to room temperature and water added. The mixture was filtered and the residue was washed with water. The residue was dried under reduced pressure. Yield: 0.43 g of compound 40 (96%).

f) Preparation of compound 70




15

Intermediate 14 (prepared according to A6) (0.001 mol) was suspended in di-(2-hydroxyethyl)-amine (1 ml), and the mixture was heated at 100°C for 30 minutes. After cooling, water was added with stirring, and the mixture was filtered. The residue was washed with water, and the residue was dried under reduced pressure.

20 Yield : 0.18 g of compound 70 (42%).


-64-

g) Preparation of compound 87




Intermediate 14a (prepared according to A6) (0.0001 mol) was dissolved in tetrahydrofuran (4 ml), and 2-aminoethanol (0.0002 mol) was added. The solution was stirred at room temperature for 1 hour, the solvent removed under reduced pressure and the residue was dissolved in MeOH (2 ml). The solution was acidified with a 6 M solution of hydrogen chloride in 2-propanol. 2-propanone (4 ml) was added, and the mixture was stirred for 16 hours. The mixture was filtered and the residue dried under reduced pressure. Yield : 0.354 g of compound 87 (87%).

h) Preparation of compound 90



Intermediate 14 (prepared according to A6) (0.0005 mol) was added to dimethylsulphoxide (5 ml) and the suspension was warmed gently until dissolution was complete. Glycine (0.001 mol) and then sodium carbonate (0.001 mol) were added. The mixture was stirred at 120°C for 4 hours, cooled to 100°C, and then water (5 ml) was added. The solution was neutralised with 1 M hydrochloric acid. Water (3 ml) was added and then cooled with rapid stirring to 0°C. The mixture was filtered and the residue dried under reduced pressure. Yield : 0.192 g of compound 90 (92%).


i) Preparation of compound 89



Ethan-1,2-diol (1 ml) was added dropwise to a mixture of sodium hydride (60% in oil, 0.0005 mol) in *N,N*-dimethylformamide, and stirring continued for 30 minutes. Intermediate 14a (0.0002 mol) was added and the reaction stirred for 20 hours. The solvent was removed under reduced pressure, taken up in water (5 ml) and neutralised with acetic acid. The mixture was gently warmed and acetonitrile was added slowly until dissolution was complete. After cooling, the mixture was filtered, and the residue was dried under reduced pressure and chromatographed on silica gel using dichloromethane:methanol 99:1 as eluent. Yield : 0.023 g of compound 89 (29%).

Example B8

Preparation of compound 81




Sodium methoxide (0.0028 mol) was added to a stirred solution of intermediate 17 (0.0028 mol) (prepared according to Ex. A8b) in methanol (20 ml), and stirring was continued for 16 hours.

In a separate flask guanidine hydrochloride (0.0028 mol) was suspended in methanol (15 ml) and sodium methoxide (0.0028 mol) was added, and stirring continued for 1 hour. This solution was then added to the first prepared solution. Stirring was continued for 16 hours. The solvent was removed under reduced pressure and 1,1-dimethoxy-*N,N*-dimethyl-methanamine (0.0028 mol), followed by a solution of sodium methoxide (0.0028 mol) in methanol (5ml), was added. The reaction was stirred for 16 hours. The solvent was removed under reduced pressure, methanol (5 ml) was added, and the solvent again removed under reduced pressure. The residue was suspended in hot ethanol, and the mixture was filtered whilst hot. The cool filtrate was filtered and the residue dried under reduced pressure. Yield : 0.080 g of compound 81 (7%).

Example B9

Preparation of compound 2




A solution of acetyl chloride (1.5 ml) in dichloromethane (10 ml) was dropwise added to a mixture of compound 32 (0.019 mol) in pyridine (30 ml). The mixture was stirred at room temperature for 75 hours, water (30 ml) was added and the mixture was filtered. The residue was washed with hexane and recrystallized from acetic acid. Yield: 3.55 g of compound 2 (54%).

-66-


Example B10

a) Preparation of compound 49




[1-(chloromethyl)-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane bis(tetrafluoroborate)] "Selectfluor" (0.003 mol) was added to a solution of compound 9 (0.0024 mol) (prepared according to B1) and 2,6-lutidine (0.045 mol) in *N,N*-dimethylformamide (5 ml). The reaction mixture was stirred at room temperature for 48 hours, and the volatile components removed under reduced pressure. The residue was chromatographed on silica gel using tetrahydrofuran:hexane 1:4 as eluent. Yield : 0.093 g of compound 49 (11%).

b) Preparation of compound 16



N-Chlorosuccinimide (0.0038 mol) was added to a solution of compound 9 (0.0035 mol) (prepared according to B1) in carbon tetrachloride (30 ml). The mixture was refluxed for 5 hours, and the solvent was removed at reduced pressure. The residue was suspended in water, the mixture boiled for 5 minutes, and the mixture filtered. The residue was recrystallised from ethanol. Yield : 0.81 g of compound 16 (62%).

c) Preparation of compound 11




N-Bromosuccinimide (0.0026 mol) was added to a solution of compound 6 (prepared according to B1) (0.0024 mol) in carbon tetrachloride (3 ml). The mixture was refluxed for 4 hours, and the solvent removed at reduced pressure. The residue was recrystallised from ethanol:water (4:1), and then from ethanol. Yield : 0.89 g of compound 11 (89%).

-67-

Example B11


Preparation of compound 42



Compound 40 (prepared according to B7e) (0.00087 mol) was dissolved in concentrated sulphuric acid (98%, 28 ml) and the mixture was heated to 40°C. Water (0.35 ml) was slowly added. After stirring for 2 hours, the reaction mixture was poured onto ice and neutralised with cold aqueous ammonia. The mixture was filtered, and the residue washed with water and then with ethanol:diethyl ether (1:5). Yield : 0.25 g of compound 42 (79%).

Example B12


Preparation of compound 80



A mixture of guanidine hydrochloride (0.025 mol) and sodium methoxide (0.025 mol) in 2-ethoxyethanol (20 ml) were refluxed for 15 minutes, and then intermediate 2a (0.013 mol) was added in one portion. Stirring was continued under reflux for 90 minutes, and the solution was cooled and diluted with ethanol and water. The pH was adjusted to 3 using acetic acid. The mixture was filtered, and the residue washed with water and then dried under reduced pressure. Yield : 3.6 g of compound 80 (79%).

Example B13

Preparation of compound 99




Sodium hydride (0.020 mol) was added to a solution of guanidine hydrochloride (0.020 mol) in *N,N*-dimethylformamide (10 ml) and stirring continued for 30 minutes. A solution of intermediate 18 (0.014 mol) in *N,N*-dimethylformamide (10 ml) was added, and the reaction was heated under reflux for 2 hours. The solution was cooled to 0 °C, water (150 ml) was added, and the mixture was filtered. The residue was dried under

reduced pressure. Yield : 1.5 g of compound 99 (28%). A sample was recrystallised from acetonitrile to give compound 99 as a yellow solid (mp. 178-180 °C).

Example B14

Preparation of compound 97




- 5 Intermediate 20 (0.001 mol) was dissolved in tetrahydrofuran (10 ml) in an autoclave and liquid ammonia (0.6 mol) was added. The autoclave was closed and the reaction stirred at room temperature for 16 hours. The mixture was filtered, and the volatile components removed from the filtrate under reduced pressure. The residue was chromatographed on silica gel using dichloromethane:methanol 99:1 as eluent. Yield :
- 10 0.15 g of compound 97 (42%).

Compounds 92, 93, 94, 95, 96 and 98 were made analogously according to the method described in Example B7a) or B7f).


- 15 Tables 1, 2 and 3 list the compounds of formula (I) which were prepared according to, analogue to one of the examples and methods described above.

Table 1



















































Comp. No.	Ex. No.	R ^a	R ^b	R ^c	R ^d	Physical data (m.p.)
1	B1	H	H	H	H	203
2	B9	-C(=O)-CH ₃	H	4-Cl	H	>280
3	B1	H	H	2-Cl	H	209
4	B1	H	H	2-Cl	3-Cl	220
5	B1	H	H	2-Cl	4-Cl	
6	B1	H	H	3-CF ₃	H	168
7	B1	H	H	3-Cl	H	175
8	B1	H	H	3-F	H	198
10	B1	H	H	4-F	H	221
11	B10c	H	5-Br	3-CF ₃	H	166
12	B10c	H	5-Br	3-F	H	136
14	B10c	H	5-Br	4-F	H	169
15	B10b	H	5-Cl	3-CF ₃	H	174
17	B3a	CH ₃	H	4-Cl	H	
32	B1	H	H	4-Cl	H	
43	B1	H	H	4-CH ₃	H	209
80	B12	H	6-OH	4-CF ₃	H	>270



























Table 2



Comp. No.	Ex.No.	Q	L	Z	Physical data (m.p.)
20	B1	2-pyridyl		CH ₃	248
21	B1	3-pyridyl		CH ₃	
22	B3b	4-chlorophenyl	4-pyrimidinyl	CH ₃	191
23	B2	phenyl	3-pyrazolyl	CH ₃	200
24	B2	4-chlorophenyl	3-pyrazolyl	CH ₃	204
25	B2	2-pyridyl	3-pyrazolyl	CH ₃	190
26	B2	3-pyridyl	3-pyrazolyl	CH ₃	177
27		phenyl		CH ₃	138
28	B2	4-pyridyl	3-pyrazolyl	CH ₃	198
29	B3a	4-pyridyl		CH ₃	188
30	B3b	4-pyridyl	4-pyrimidinyl	CH ₃	144
31	B1	4-pyridyl		CH ₃	249
32	B1	1-pyrrolyl		CH ₃	188
33	B1	2-thienyl		CH ₃	198
34	B1	1-pyrrolyl		Cl	239
44	B1	4-fluorophenyl		Cl	241
45	B1	phenyl		Cl	213

Comp. No.	Ex.No.	Q	L	Z	Physical data (m.p.)
46	B1			CH ₃	197
47	B1			CH ₃	251
48	B1			CH ₃	247
50	B1			Cl	240
36	B5			C(=O)-NH ₂	287
51	B3c			CH ₃	
52	B1			OCH ₃	177
53	B3a			CH ₃	.HCl (1:1) 194
54	B3a			Cl	220
55	B3c			CH ₃	172
56	B3a			CH ₃	.HCl(1:2) .H ₂ O(1:1) > 250
57	B3c			Cl	184
58	B3c			Cl	177
59	B3c			Cl	182

Comp. No.	Ex.No.	Q	L	Z	Physical data (m.p.)
67	B7a			CH ₃	184
68	B7a			CH ₃	139
69	B7a			CH ₃	124
70	B7f			CH ₃	164-180
35	B4			CH ₃	106
42	B11			CH ₃	231
40	B7e			CH ₃	214
76	B3a			CH ₃	188
77	B7a			CH ₃	114
78	B3a			CH ₃	

Comp. No.	Ex.No.	Q	L	Z	Physical data (m.p.)
37	B6			CH ₃	123
79	B6			CH ₃	
81	B8			CH ₃	208
82	B7d			CH ₃	>260
83	B7c			CH ₃	128
86	B3a			Cl	216
87	B7g			Cl	HCl 210
88	B7a			Cl	>250
89	B7i			Cl	152
91	B6			Cl	146
92	B7f			CH ₃	174
93	B7f			CH ₃	145 (decomposition)
94	B7f			CH ₃	HCl 210 (decomposition)














Comp. No.	Ex.No.	Q	L	Z	Physical data (m.p.)
95	B7a			CH ₃	84 (decomposition)
96	B7a			CH ₃	HCl 260 (decomposition)
97	B14			CH ₃	207
98	B7c			CH ₃	163
99	B13			CH ₃	

Table 3



Comp. No.	Ex. No.	R ^a	R ^b	Physical data (m.p.)
9	B1	H	H	186
13	B10c	H	5-Br	190
16	B10b	H	5-Cl	186
49	B10a	H	5-F	181
60	B7a	-CH ₂ -CH ₂ -NH ₂	H	171
38	B7a	-CH ₂ -CH ₂ -CH ₂ -NH ₂	H	139
61	B7a	-(CH ₂) ₆ -NH ₂	H	120-130
62	B7a	-(CH ₂) ₈ -NH ₂	H	110-120
63	B7g	-CH ₂ -CH ₂ -OH	H	188
64	B7g	-CH ₂ -CH ₂ -CH ₂ -OH	H	173
65	B7g	-CH ₂ -CH(OH)-CH ₂ -OH	H	194

Comp. No.	Ex. No.	R ^a	R ^b	Physical data (m.p.)
66	B7a	-C(=NH)-NH ₂	H	241
71	B7a	-(CH ₂) ₇ -NH ₂	H	110
72	B7a		H	185
73	B7a		H	155
74	B7g	-CH(CH ₂ -OH)-CH ₂ -OH	H	201
75	B7a	-(CH ₂) ₂ -O-(CH ₂) ₂ -O-(CH ₂) ₂ -NH ₂	H	89
84	B7b	-CH ₂ -CH ₂ -C(=O)O ⁻ Na ⁺	H	>240
85	B7b	-CH ₂ -C(=O)O ⁻ Na ⁺	H	
90	B7h	-CH ₂ -COOH	H	248

C. Pharmacological example

Example C.1 : *in vitro* inhibition of TNF- α production in human blood

Human whole blood stimulation

- 5 Peripheral blood from healthy male donors was drawn into heparinized syringes (12.5 U heparin/ml). Blood samples were three-fold diluted in RPMI 1640 medium (Life Technologies, Belgium) supplemented with 2 mM L-glutamine, 100 U/ml penicillin and 100 μ g/ml streptomycin, and 300 μ l fractions were distributed in 24-well multidisc plates (Nunc, Roskilde, Denmark). Blood samples were preincubated (60
- 10 minutes at 37°C) in a humidified 6% CO₂-atmosphere with 100 μ l of drug solvent (final concentration 0.02% dimethylsulfoxide in RPMI 1640) or with 100 μ l of an appropriate dose of test compound before being stimulated by the addition of 100 μ l of lipopolysaccharide at a final concentration of 100 ng/ml. After 6 hours, cell-free supernatant fluids were collected by centrifugation and stored at -20°C until tested for
- 15 the presence of TNF- α .

Example C.2 : *in vitro* inhibition of IL-12p40 production in human blood

Human whole blood stimulation

- Peripheral blood from healthy male donors was drawn into heparinized syringes (12.5 U heparin/ml). Blood samples were three-fold diluted in RPMI 1640 medium (Life Technologies, Belgium) supplemented with 2 mM L-glutamine, 100 U/ml penicillin and 100 μ g/ml streptomycin, and 300 μ l fractions were distributed in 24-well multidisc plates (Nunc, Roskilde, Denmark). Blood samples were preincubated (60
- 20 minutes at 37°C) in a humidified 6% CO₂-atmosphere with 100 μ l of drug solvent (final concentration 0.02% dimethylsulfoxide in RPMI 1640) or with 100 μ l of an
- 25

appropriate dose of test compound before being stimulated by the addition of 100 μ l of lipopolysaccharide at a final concentration of 100 ng/ml. After 24 hours, cell-free supernatant fluids were collected by centrifugation and stored at -20°C until tested for the presence of IL-12 p40.

5

Example C.3 : cytokine measurements

Cytokine protein concentrations were determined by sandwich ELISA as described in Van Wauwe et al. (1996, *Inflamm Res*, 45, 357-363). Murine monoclonals used as capture antibodies to human cytokines were obtained from R&D Systems (Abingdon, United Kingdom) and code named MAB210 and MAB611 for TNF- α and IL-12 p40 respectively. Biotinylated goat polyclonal antibodies used to detect human cytokines were from R&D Systems (BAF210, BAF219). Cytokine levels were calculated from standard curves using recombinant cytokines supplied by R&D Systems.

10

15 Example C.4 : *in vitro* inhibition of IL-12p70 production in human blood

Human whole blood stimulation

Peripheral blood from healthy donors was drawn into heparinized syringes (12.5 U heparin/ml). Blood samples were three-fold diluted in RPMI 1640 medium (Life Technologies, Belgium) supplemented with 2 mM L-glutamine, 100 U/ml penicillin and 100 μ g/ml streptomycin, and 200 μ l fractions were distributed in 96-well plates (Nunc, Roskilde, Denmark). Blood samples were preincubated (5 minutes at 37°C) in a humidified 5% CO₂-atmosphere with 25 μ l of drug solvent (final concentration 0.1 % dimethylsulfoxide in RPMI 1640) or with 25 μ l of an appropriate dose of test compound before being stimulated by the addition of 25 μ l of lipopolysaccharide at a final concentration of 100 ng/ml. After 24 hours, cell-free supernatant fluids were collected by centrifugation and stored at -20°C until tested for the presence of IL-12.

20

25

Example C.5 : cytokine measurements

Cytokine protein concentrations were determined by sandwich ELISA as described in Van Wauwe et al. (1996, *Inflamm Res*, 45, 357-363). The quantikine HS kit (R &D HS120, Abingdon, United Kingdom) was used to quantify the cytokine levels (IL12 p70) in the supernatant.

30

Table 4 lists the percentage inhibition of TNF- α and IL-12 production (column “%inhib.”) at a test dose of 1×10^{-6} , 1×10^{-7} or 1×10^{-8} M for the compounds of the present invention.

35


-77-

Table 4

Comp. No	% inhib. TNF- α		% inhib. IL-12 (p40)	% inhib. IL-12 (p70)
	1×10^{-6} M	1×10^{-7} M	1×10^{-6} M	1×10^{-8} M
1	50	51	55	
5			46	
8			59	
9			57	92
10			53	
20	54		51	
32	55	54	56	
34				95
35				94
38				93
40				93
49				92
58				84
63				93
70				90
73				92
74				82
80			49	
81				94
83			59	
84				95
90			56	


Claims

1. Use of a compound for the manufacture of a medicament for the prevention or the treatment of diseases mediated through TNF- α (Tumor Necrosis Factor-alpha) and/or IL-12 (Interleukin 12), wherein the compound is a compound of formula



a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof,
wherein

- 10 Z is halo; C₁₋₆alkyl; C₁₋₆alkyl substituted with hydroxy, carboxyl, cyano, amino, mono - or di(C₁₋₆alkyl)amino, aminocarbonyl, mono -or di(C₁₋₆alkyl)aminocarbonyl, C₁₋₆alkyloxycarbonyl or C₁₋₆alkyloxy; polyhaloC₁₋₄alkyl; C₁₋₄alkyloxy; cyano; amino; aminocarbonyl; mono -or di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; H₂N-S(=O)₂-; mono -or di(C₁₋₆alkyl)amino-S(=O)₂; -C(=N-R^x)NR^yR^z;
- 15 R^x is hydrogen, C₁₋₆alkyl, cyano, nitro or -S(=O)₂-NH₂;
R^y is hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl or C₂₋₆alkynyl;
R^z is hydrogen or C₁₋₆alkyl;
- 20 Q is C₃₋₆cycloalkyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, phenyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl, or imidazopyridyl, wherein each of said ring systems may optionally be substituted with up to three substituents each of said substituents independently being selected from halo; hydroxy; cyano; carboxyl; azido; amino; mono- or di(C₁₋₆alkyl)amino; C₁₋₆alkylcarbonylamino; C₁₋₆alkyl;
- 25 C₂₋₆alkenyl; C₂₋₆alkynyl; C₃₋₆cycloalkyl; C₁₋₆alkyl substituted with hydroxy, C₁₋₆alkyloxy, amino, mono-or di(C₁₋₄alkyl)amino; C₁₋₆alkyloxy; C₁₋₆alkylthio; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; arylC₁₋₆alkyloxy; aryloxy; polyhaloC₁₋₆alkyl; polyhaloC₁₋₆alkyloxy; polyhaloC₁₋₆alkylcarbonyl; C₁₋₄alkyl-S(=O)_n- or R¹HN-S(=O)_n-, with R¹ representing hydrogen, or a radical
- 30 of formula




with A being O, S or a bivalent radical of formula -CR²=N- with CR² attached to N of formula (a-1); and

-79-

R^2 being hydrogen, C_{1-6} alkyl or C_{1-6} alkyloxy;

or

Q is a radical of formula



5 wherein B_1 and B_2 each independently are O, NR^3 , CH_2 or S, with R^3 being hydrogen or C_{1-4} alkyl;

B_3 is O or NR^4 with R^4 being hydrogen or C_{1-4} alkyl;

q is an integer with value 1 to 4;

r is an integer with value 1 to 3;

10 n is an integer with value 1 or 2;

L is phenyl substituted with up to 4 substituents each substituent independently being selected from C_{1-6} alkyloxycarbonyl; C_{1-6} alkylcarbonyloxy; aminocarbonyl; mono- or di(C_{1-6} alkyl)aminocarbonyl; C_{1-6} alkyl-C(=O)-NH-; C_{1-6} alkyloxy-C(=O)-NH-; $H_2N-C(=O)-NH-$; mono- or di(C_{1-4} alkyl)amino-C(=O)-NH-; Het-NH-; Het¹-NH-; -NH-C(=N- R^x) NR^yR^z ; -C(=N- R^x) NR^yR^z ; Het¹; or a radical of formula

15 -X-C₁-Y₁-C₂-Y₂-C₃-Y₃-C₄-Z (c-1) wherein

X represents NR^5 , O, S or a direct bond;

C₁, C₂, C₃ and C₄ each independently represent C_{1-6} alkanediyl, C_{2-6} alkenediyl, C_{2-6} alkynediyl or a direct bond;

20 Y₁, Y₂ and Y₃ each independently represent NR^5 , O, S or a direct bond;

Z is hydrogen, halo, cyano, hydroxy, carboxyl, -P(=O)(OH)H,

-P(=O)(OH)₂, -P(=O)(OH)CH₃, -P(=O)(OH)(OCH₃),

-P(=O)(OH)(OCH₂CH₃), -P(=O)(OH)NH₂, -S(=O)₂H, -S(=O)₂(OH),

25 -S(=O)₂NH, -C(=O)-NH-S(=O)₂-H, tetrazolyl, 3-hydroxy-isothiazolyl, 3-hydroxy-isoxazolyl, 3-hydroxy-thiadiazolyl, mercaptotetrazolyl,

3-mercapto-triazolyl, 3-sulfinyl-triazolyl, 3-sulfonyl-triazolyl;

R^5 is hydrogen, C_{1-6} alkyl or -C(=NH)-N(R^z)₂; and wherein

from 1 to 3 hydrogen atoms of the C_{1-6} alkyl, C_{1-6} alkanediyl, C_{2-6} alkenediyl or C_{2-6} alkynediyl groups in the definitions of R^5 and the radical of formula (c-1) may optionally and each independently be

30 replaced by halo, hydroxy, carboxyl, -P(=O)(OH)H, -P(=O)(OH)₂, -P(=O)(OH)CH₃, -P(=O)(OH)-(OCH₃), -P(=O)(OH)(OCH₂CH₃), -P(=O)(OH)NH₂, -S(=O)₂H, -S(=O)₂(OH), -S(=O)₂NH, -C(=O)-NH-S(=O)₂-H, tetrazolyl, 3-hydroxy-isothiazolyl,

-80-

3-hydroxy-isoxazolyl, 3-hydroxy-thiadiazolyl, mercaptotetrazolyl,
3-mercapto-triazolyl, 3-sulfinyl-triazolyl, 3-sulfonyl-triazolyl;

or

L is a monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a
bicyclic partially saturated or aromatic heterocycle wherein each of said ring
systems may optionally be substituted with up to 3 substituents, each substituent
independently being selected from C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy;
aminocarbonyl; mono-or di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyl-C(=O)-NH-;
C₁₋₆alkyloxy-C(=O)-NH-; H₂N-C(=O)-NH-; mono- or di(C₁₋₄alkyl)amino-C(=O)-
NH-; Het-NH-; Het¹-NH-; -NH-C(=N-R^x)NR^yR^z; -C(=N-R^x)NR^yR^z; Het¹; or a
radical of formula

-X-C₁-Y₁-C₂-Y₂-C₃-Y₃-C₄-Z (c-1) wherein

X represents NR⁵, O, S or a direct bond;
C₁, C₂, C₃ and C₄ each independently represent C₁₋₆alkanediyl,
C₂₋₆alkenediyl, C₂₋₆alkynediyl or a direct bond;
Y₁, Y₂ and Y₃ each independently represent NR⁵, O, S or a direct bond;
Z is hydrogen, halo, cyano, hydroxy, carboxyl, -P(=O)(OH)H,
-P(=O)(OH)₂, -P(=O)(OH)CH₃, -P(=O)(OH)(OCH₃),
-P(=O)(OH)(OCH₂CH₃), -P(=O)(OH)NH₂, -S(=O)₂H, -S(=O)₂(OH),
-S(=O)₂NH, -C(=O)-NH-S(=O)₂-H, tetrazolyl, 3-hydroxy-isothiazolyl,
3-hydroxy-isoxazolyl, 3-hydroxy-thiadiazolyl, mercaptotetrazolyl,
3-mercapto-triazolyl, 3-sulfinyl-triazolyl, 3-sulfonyl-triazolyl;
R⁵ is hydrogen, C₁₋₆alkyl or -C(=NH)-N(R^z)₂; and wherein
from 1 to 3 hydrogen atoms of the C₁₋₆alkyl, C₁₋₆alkanediyl,
C₂₋₆alkenediyl or C₂₋₆alkynediyl groups in the definitions of R⁵ and the
radical of formula (c-1) may optionally and each independently be
replaced by halo, hydroxy, carboxyl, -P(=O)(OH)H, -P(=O)(OH)₂,
-P(=O)(OH)CH₃, -P(=O)(OH)-(OCH₃), -P(=O)(OH)(OCH₂CH₃),
-P(=O)(OH)NH₂, -S(=O)₂H, -S(=O)₂(OH), -S(=O)₂NH,
-C(=O)-NH-S(=O)₂-H, tetrazolyl, 3-hydroxy-isothiazolyl,
3-hydroxy-isoxazolyl, 3-hydroxy-thiadiazolyl, mercaptotetrazolyl,
3-mercapto-triazolyl, 3-sulfinyl-triazolyl, 3-sulfonyl-triazolyl;

Het is a monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a
bicyclic partially saturated or aromatic heterocycle wherein each of said ring
systems may optionally be substituted with up to 3 substituents, each substituent
independently being selected from halo, hydroxy, amino, cyano, carboxyl,
mono-or di(C₁₋₆alkyl)amino, C₁₋₆alkyl, C₁₋₆alkyl substituted with hydroxy or

C₁₋₄alkyloxy or amino or mono-or di(C₁₋₄alkyl)amino, polyhaloC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylcarbonyloxy, aminocarbonyl, mono-or di(C₁₋₆alkyl)aminocarbonyl, C₁₋₆alkyl-C(=O)-NH-, C₁₋₆alkyloxy-C(=O)-NH-, H₂N-C(=O)-NH- or mono- or di(C₁₋₄alkyl)amino-C(=O)-NH-;


Het¹ is a saturated 6-membered heterocycle selected from piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, wherein said saturated 6-membered heterocycle may optionally be substituted with amino or C₁₋₄alkyl optionally substituted with aryl;

aryl is phenyl, optionally substituted with up to five substituents each independently selected from halo, hydroxy, C₁₋₆alkyl, polyhaloC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, cyano, nitro, amino, mono-or di(C₁₋₆alkyl)amino.

2. Use of a compound as claimed in claim 1 wherein Q is C₃₋₆cycloalkyl, phenyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl, or imidazopyridyl, each of said rings optionally being substituted with up to three substituents each independently selected from halo; hydroxy; cyano; carboxyl; azido; amino; mono- or di(C₁₋₆alkyl)amino; C₁₋₆alkylcarbonylamino; C₁₋₆alkyl; C₂₋₆alkenyl; C₂₋₆alkynyl; C₃₋₆cycloalkyl; C₁₋₆alkyl substituted with hydroxy, C₁₋₆alkyloxy, amino, mono-or di(C₁₋₄alkyl)amino; C₁₋₆alkyloxy; C₁₋₆alkylthio; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; arylC₁₋₆alkyloxy; aryloxy; polyhaloC₁₋₆alkyl; polyhalo-C₁₋₆alkyloxy; polyhaloC₁₋₆alkylcarbonyl; C₁₋₄alkyl-S(=O)_n- or R¹HN-S(=O)_n-;

or

Q is a radical of formula



wherein X and Y each independently are O, NR³, CH₂ or S, with R³ being hydrogen or C₁₋₄alkyl;

q is an integer with value 1 to 4;

Z is O or NR⁴ with R⁴ being hydrogen or C₁₋₄alkyl;

r is an integer with value 1 to 3;

Z is halo; C₁₋₆alkyl; C₁₋₆alkyl substituted with hydroxy, carboxyl, cyano, amino, mono- or di(C₁₋₆alkyl)amino, aminocarbonyl, mono- or di(C₁₋₆alkyl)aminocarbonyl, C₁₋₆alkyloxycarbonyl, C₁₋₆alkyloxy; polyhaloC₁₋₄alkyl; cyano; amino; aminocarbonyl;

-82-

mono -or di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; aminoS(=O)₂-; mono -or di(C₁₋₆alkyl)aminoS(=O)₂; -C(=N-R^x)NR^yR^z;

L is phenyl, substituted with up to 4 substituents each independently being selected from halo, hydroxy, mercapto, amino, cyano, carboxyl, mono-or
 5 di(C₁₋₆alkyl)amino, C₁₋₆alkyl, C₁₋₆alkyl substituted with hydroxy or C₁₋₄alkyloxy or amino or mono-or di(C₁₋₄alkyl)amino, polyhaloC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylcarbonyloxy, aminocarbonyl, mono-or di(C₁₋₆alkyl)aminocarbonyl, C₁₋₆alkyl-C(=O)-NH-, C₁₋₆alkyloxy-C(=O)-NH-, H₂N-C(=O)-NH-, mono- or di(C₁₋₄alkyl)amino-
 10 C(=O)-NH- or Het-NH-, -C(=N-R^x)NR^yR^z; or

L is a monocyclic 5 or 6-membered partially saturated or aromatic heterocycle or a bicyclic partially saturated or aromatic heterocycle wherein each of said ring systems may optionally be substituted with up to 3 substituents, each substituent
 15 independently being selected from halo, hydroxy, mercapto, amino, cyano, carboxyl, mono-or di(C₁₋₆alkyl)amino, C₁₋₆alkyl, C₁₋₆alkyl substituted with hydroxy or C₁₋₄alkyloxy or amino or mono-or di(C₁₋₄alkyl)amino, polyhaloC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylcarbonyloxy, aminocarbonyl, mono-or di(C₁₋₆alkyl)aminocarbonyl, C₁₋₆alkyl-C(=O)-NH-, C₁₋₆alkyloxy-C(=O)-NH-, H₂N-C(=O)-NH-, mono- or
 20 di(C₁₋₄alkyl)amino-C(=O)-NH- or Het-NH-, -C(=N-R^x)NR^yR^z.

3. Use of a compound as claimed in claim 1 or 2 wherein L is furanyl, pyrrolyl, oxazolyl, imidazolyl, pyrazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, a
 25 5-membered partially saturated heterocycle, a 6-membered partially saturated or aromatic heterocycle or a bicyclic partially saturated or aromatic heterocycle wherein each of the aforesaid ring systems may optionally be substituted with up to 3 substituents, each substituent independently being selected from C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; aminocarbonyl; mono-or di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyl-C(=O)-NH-; C₁₋₆alkyloxy-C(=O)-NH-; H₂N-C(=O)-NH-; mono- or di(C₁₋₄alkyl)amino-
 30 C(=O)-NH-; Het-NH-; Het¹-NH-; -NH-C(=N-R^x)NR^yR^z; -C(=N-R^x)NR^yR^z; Het¹; or a radical of formula -X-C₁-Y₁-C₂-Y₂-C₃-Y₃-C₄-Z provided L is other than optionally substituted quinoxaliny.

4. Use of a compound as claimed in any one of claims 1 to 3 wherein L is a
 35 6-membered partially saturated or aromatic heterocycle wherein each of said ring systems may optionally be substituted with up to 3 substituents, each substituent independently being selected from C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy;

aminocarbonyl; mono-or di(C₁₋₆alkyl)aminocarbonyl; C₁₋₆alkyl-C(=O)-NH-;
 C₁₋₆alkyloxy-C(=O)-NH-; H₂N-C(=O)-NH-; mono- or di(C₁₋₄alkyl)amino-C(=O)-NH-;
 Het-NH-; Het¹-NH-; -NH-C(=N-R^x)NR^yR^z; -C(=N-R^x)NR^yR^z; Het¹; or a radical of
 formula -X-C₁-Y₁-C₂-Y₂-C₃-Y₃-C₄-Z.












5





5. Use of a compound as claimed in claim 1 wherein Q is phenyl, pyridyl, pyrrolyl,
 pyrazolyl or thienyl, wherein each of said ring systems may optionally be substituted
 with one or two substituents each independently being selected from halo or
 polyhaloC₁₋₆alkyl; Z is C₁₋₆alkyl, halo, C₁₋₆alkyloxy, aminocarbonyl; L is pyrimidinyl,
 10 pyrazolyl, triazolyl or triazinyl, wherein each of said ring systems may optionally be
 substituted with one or two substituents each independently being selected from amino,
 C₁₋₆alkylcarbonylamino, halo, Het-NH-, hydroxy, C₁₋₆alkylthio, C₁₋₆alkyloxy, C₁₋₆alkyl,
 C₁₋₁₂alkylamino, mono-or di(hydroxyC₁₋₁₂alkyl)amino wherein C₁₋₁₂alkyl may further
 optionally be substituted with hydroxy, Het¹, aminocarbonyl, cyano,
 15 aminoC₁₋₁₂alkylamino, hydroxyC₁₋₁₂alkyloxy, -NH-C(=NH)-NH₂,
 carboxyC₁₋₁₂alkylamino or aminoC₁₋₆alkyloxyC₁₋₆alkyloxyC₁₋₆alkylamino..

6. A compound as defined in any one of claims 1 to 5 provided that
 -L is other than substituted phenyl;
 20 - when Z is methyl, Q is phenyl or phenyl substituted with halo, methyl or ethyloxy,
 then L is other than quinoxalinyll;
 - when Z is methyl, Q is phenyl or phenyl substituted at the para position with methyl,
 chloro, nitro or methyloxy, then L is other than thiazolyl substituted with methyl or
 amino;
 25 - when Z is trifluoromethyl, Q is 4-methylphenyl, then L is other than 1,2,3-triazolyl
 mono-or disubstituted with methyloxycarbonyl;
 - L is other than unsubstituted or substituted benzoxazolyl or unsubstituted or
 substituted benzimidazolyl;

-84-







- the compound is other than

L	Z	Q
	CH ₃ -	phenyl
3-pyridyl	CH ₃ -	4-pyridyl
	CH ₃ -C(=O)-O-	phenyl
	CH ₃ -	4-chloro-phenyl
	CH ₃ -	phenyl
	-CF ₃	4-methyl-phenyl
	-CH ₂ -C(=O)-O-CH ₂ CH ₃	4-chloro-phenyl
	-CF ₃	phenyl
	CH ₃ -	phenyl
	CH ₃ -	4-chlorophenyl
	CH ₃ -	phenyl
	CH ₃ -	2-thienyl

	F-	
	CH ₃ -	phenyl
	CH ₃ -	4-chlorophenyl

7. A compound as defined in any one of claims 1 to 5 provided that
- L is other than substituted phenyl;
 - when Z is methyl, Q is phenyl or phenyl substituted with halo, CH₃ or ethyloxy, then L
- 5 is other than quinoxaliny;
- when Z is methyl, Q is phenyl or phenyl substituted at the para position with methyl, chloro, nitro or methyloxy, then L is other than thiazolyl substituted with methyl or amino;
 - the compound is other than

10

L	Z	Q
3-pyridyl	CH ₃ -	4-pyridyl
	CH ₃ -	4-chloro-phenyl
	CH ₃ -	phenyl
	-CH ₂ -C(=O)-O-CH ₂ CH ₃	4-chloro-phenyl
	CH ₃ -	phenyl
	F-	


for use as a medicine.

-86-

8. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and as active ingredient a therapeutically effective amount of a compound as claimed in claim 6 or 7.

5 9. A process of preparing a composition as claimed in claim 8 characterized in that a pharmaceutically acceptable carrier is intimately mixed with a therapeutically effective amount of a compound as claimed in claim 6 or 7.


10. A compound of formula



10

wherein Z^b represents halo or C_{1-4} alkyl.


11. A compound of formula



15

wherein Z^b represents halo or C_{1-4} alkyl.

12. A process of preparing a compound as claimed in claim 6 characterized by
a) reacting an intermediate of formula (II) with an intermediate of formula (III) or a suitable salt thereof, in the presence of a suitable solvent and a suitable alcoholate




20

with Q and Z as defined in claim 1 and R^a representing hydrogen, amino, optionally substituted C_{1-6} alkyl, optionally substituted mono- or di(C_{1-12} alkyl)amino, Het-NH- or Het¹;


b) reacting an intermediate of formula (II') with an intermediate of formula (III) or a
25 suitable salt thereof, in the presence of a suitable solvent and a suitable alcoholate,

-87-



with Q as defined in claim 1 and R^a representing hydrogen, amino, optionally substituted C_{1-6} alkyl, optionally substituted mono-or di(C_{1-12} alkyl)amino, Het-NH- or Het¹;


- 5 c) reacting an intermediate of formula (IV) or (IV') with an intermediate of formula (V) optionally at elevated temperature and optionally in the presence of a suitable solvent, and optionally in the presence of a suitable base or a suitable acid, and optionally under pressure



10 with Q and Z as defined in claim 1, R^a representing hydrogen, amino, optionally substituted C_{1-6} alkyl, optionally substituted mono-or di(C_{1-12} alkyl)amino, Het-NH- or Het¹, and R^b representing $-NH_2$, Het¹-NH-; Het¹; $-NH-C(=NH)-N(R^2)_2$; C_{1-12} alkyloxy optionally substituted with one, two or three hydroxy groups; optionally substituted

15 mono-or di(C_{1-12} alkyl)amino, in particular unsubstituted mono-or di(C_{1-12} alkyl)amino or mono-or di(C_{1-12} alkyl)amino wherein C_{1-12} alkyl is substituted with one, two or three substituents selected from hydroxy, carboxyl, amino, amino C_{1-4} alkyloxy C_{1-4} alkyloxy;


d) reacting an intermediate of formula (IV) with a suitable cyanide salt in the presence of a suitable solvent



-88-


with Q and Z as defined in claim 1;

e) reacting an intermediate of formula (IV) with a suitable hydroxyde base in the presence of a suitable solvent




5 with Q and Z as defined in claim 1;

f) reacting an intermediate of formula (II) with thiourea in the presence of a suitable solvent, a suitable alcoholate, dimethyl sulphate, and a suitable base




with Q and Z as defined in claim 1;

10 g) reacting an intermediate of formula (XXXVI) with an intermediate of formula (III) in the presence of a suitable solvent



with Q and Z as defined in claim 1 and R^a as defined above in process variant a) and b);


15 h) reacting an intermediate of formula (II) with hydrazine monohydrate in the presence of a suitable acid



with Q and Z as defined in claim 1;


i) reacting an intermediate of formula (VII) with an intermediate of formula (VIII) in the presence of a suitable base and a suitable solvent

-89-




with Q and Z as defined in claim 1 and R^c representing hydrogen or C₁₋₆alkyl;

j) reacting an intermediate of formula (VI) with an intermediate of formula (III) or a suitable salt thereof, in the presence of a suitable solvent



with Q and Z as defined in claim 1 and R^a as defined above in process variant a) and b);

k) reacting an intermediate of formula (XXX) with an intermediate of formula (III) or a suitable salt thereof, in the presence of 1,1-dimethoxy-N,N-dimethyl-methanamine and in the presence of a suitable solvent



with Q and Z as defined in claim 1 and R^a as defined above in process variant a) and b);

and, if desired, converting compounds of formula (I) into each other following art-known transformations, and further, if desired, converting the compounds of formula (I), into a therapeutically active non-toxic acid addition salt by treatment with an acid, or into a therapeutically active non-toxic base addition salt by treatment with a base, or conversely, converting the acid addition salt form into the free base by treatment with alkali, or converting the base addition salt into the free

acid by treatment with acid; and, if desired, preparing stereochemically isomeric forms, quaternary amines or *N*-oxide forms thereof.

- 5 13. A product containing (a) a compound as defined in claim 6 or 7, and (b) another anti-inflammatory or immunosuppressive compound, as a combined preparation for simultaneous, separate or sequential use in the treatment of inflammatory or autoimmune diseases.
- 10 14. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and as active ingredients (a) a compound as defined in claim 6 or 7, and (b) another anti-inflammatory or immunosuppressive compound.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 02/08956

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/426 A61K31/427 A61K31/506 A61K31/497 C07D417/04
C07D417/14 A61P31/12 A61P37/02 A61P37/08 A61P29/00
C07D277/24

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 01449 A (ASTRA PHARMA PROD ;PERRY MATTHEW (GB); THORNE PHILIP (GB); BANTICK) 15 January 1998 (1998-01-15) cited in the application page 13, line 27 -page 14, line 28	1-11
X	WO 96 03392 A (SEARLE & CO ;TALLEY JOHN J (US); CARTER JEFFERY S (US); COLLINS PA) 8 February 1996 (1996-02-08) cited in the application page 63, line 6-11; claims 1-38	1-14
Y	US 5 240 929 A (CONNOR DAVID T ET AL) 31 August 1993 (1993-08-31) cited in the application abstract	1-11

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

20 November 2002

Date of mailing of the international search report

29/11/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Herrera, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/08956

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 00 35911 A (RHONE-POULENC RORER LTD ; HALLEY FRANK (GB); COLLIS ALAN JOHN (GB);) 22 June 2000 (2000-06-22) cited in the application abstract page 13, line 20-24	1-14
X	I. SIMITI ET AL: "Synthese und antibakterielle Wirkung von "-Aryl-4-R-5-glyoxyloyl-thiazolen" ARCH. PHARM., vol. 314, no. 9, 1981, pages 744-750, XP001026823 Weinheim, Germany cited in the application the whole document	1-14
X	S.N. SAWHNEY ET AL: "Thiazole Derivatives: part I - Synthesis & Anti-inflammatory Activity of Some 2'-Alkyl/Aryl-2-aryl-4-methyl-4',5-bithiaz- olyls & 2'-Amino/Substituted amino-2-aryl-4-methyl-4',5-bithiazolyls" INDIAN J CHEMISTRY, vol. 14b, no. 7, 1976, pages 552-555, XP001026830 India cited in the application the whole document	1-14
P,X	WO 02 50048 A (SIERRA MICHAEL LAWRENCE ; BESWICK PAUL JOHN (GB); GLAXO GROUP LTD () 27 June 2002 (2002-06-27) claims	1-9, 12-14
P,X	page 16	10,11
P,X	WO 02 50047 A (SIERRA MICHAEL LAWRENCE ; GLAXO GROUP LTD (GB)) 27 June 2002 (2002-06-27) claims	1-14
P,Y	WO 02 059098 A (GLAXO GROUP LTD ; LAMBERT MILLARD HURST III (US); SZNAIDMAN MARCOS) 1 August 2002 (2002-08-01) claims	1-14
E	WO 02 062774 A (SIERRA MICHAEL LAWRENCE ; GOSMINI ROMAIN LUC MARIE (FR); GLAXO GROU) 15 August 2002 (2002-08-15) claims	1-14

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 02/08956

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Present claims relate to an extremely large number of possible compounds and uses. Support within the meaning of Article 84 EPC and/or disclosure within the meaning of Article 83 EPC is to be found, however, for only a very small proportion of the compounds and uses claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds specifically disclosed in the application as well as the general idea underlying the invention.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/08956

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9801449	A	15-01-1998	AU 708882 B2	12-08-1999
			AU 3711697 A	02-02-1998
			BR 9710215 A	10-08-1999
			CA 2260057 A1	15-01-1998
			CZ 9900036 A3	16-06-1999
			EE 9900012 A	15-06-1999
			EP 0915879 A1	19-05-1999
			JP 2000516206 T	05-12-2000
			PL 331090 A1	21-06-1999
			WO 9801449 A1	15-01-1998
			TR 9900013 T2	21-04-1999
WO 9603392	A	08-02-1996	AU 3201095 A	22-02-1996
			CA 2195847 A1	08-02-1996
			EP 1125932 A2	22-08-2001
			EP 0772606 A1	14-05-1997
			JP 10504542 T	06-05-1998
			WO 9603392 A1	08-02-1996
			US 5668161 A	16-09-1997
US 5240929	A	31-08-1993	AU 672619 B2	10-10-1996
			AU 4787093 A	03-03-1994
			EP 0652874 A1	17-05-1995
			JP 3223190 B2	29-10-2001
			JP 8500101 T	09-01-1996
			KR 275428 B1	15-12-2000
			US 5312821 A	17-05-1994
			WO 9403448 A1	17-02-1994
WO 0035911	A	22-06-2000	AT 227719 T	15-11-2002
			AU 1871100 A	03-07-2000
			EP 1140916 A1	10-10-2001
			WO 0035911 A1	22-06-2000
			JP 2002532495 T	02-10-2002
WO 0250048	A	27-06-2002	AU 2966902 A	01-07-2002
			WO 0250048 A1	27-06-2002
WO 0250047	A	27-06-2002	AU 3327102 A	01-07-2002
			WO 0250047 A1	27-06-2002
WO 02059098	A	01-08-2002	WO 02059098 A1	01-08-2002
WO 02062774	A	15-08-2002	WO 02062774 A1	15-08-2002

Form PCT/ISA/210 (patent family annex) (July 1992)

